

Author's Accepted Manuscript

Partial Nephrectomy is Associated with Higher Risk of Relapse Compared with Radical Nephrectomy for Clinical Stage T1 Renal Cell Carcinoma Pathologically Upstaged to T3a

Paras H. Shah , Daniel M. Moreira , Vinay R. Patel , Geoffrey Gaunay , Arvin K. George , Manaf Alom , Zachary Kozel , Oksana Yaskiv , Simon J. Hall , Michael J. Schwartz , Manish A. Vira , Lee Richstone , Louis R. Kavoussi

PII: S0022-5347(17)35379-X
DOI: [10.1016/j.juro.2017.03.012](https://doi.org/10.1016/j.juro.2017.03.012)
Reference: JURO 14599

To appear in: *The Journal of Urology*
Accepted Date: 1 March 2017

Please cite this article as: Shah PH, Moreira DM, Patel VR, Gaunay G, George AK, Alom M, Kozel Z, Yaskiv O, Hall SJ, Schwartz MJ, Vira MA, Richstone L, Kavoussi LR, Partial Nephrectomy is Associated with Higher Risk of Relapse Compared with Radical Nephrectomy for Clinical Stage T1 Renal Cell Carcinoma Pathologically Upstaged to T3a, *The Journal of Urology*® (2017), doi: 10.1016/j.juro.2017.03.012.

DISCLAIMER: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our subscribers we are providing this early version of the article. The paper will be copy edited and typeset, and proof will be reviewed before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to The Journal pertain.

Embargo Policy

All article content is under embargo until uncorrected proof of the article becomes available online.

We will provide journalists and editors with full-text copies of the articles in question prior to the embargo date so that stories can be adequately researched and written. The standard embargo time is 12:01 AM ET on that date. Questions regarding embargo should be directed to jumedia@elsevier.com.



PARTIAL NEPHRECTOMY IS ASSOCIATED WITH HIGHER RISK OF RELAPSE COMPARED WITH RADICAL NEPHRECTOMY FOR CLINICAL STAGE T1 RENAL CELL CARCINOMA PATHOLOGICALLY UPSTAGED TO T3a

Paras H. Shah¹, Daniel M. Moreira², Vinay R. Patel¹, Geoffrey Gaunay¹, Arvin K. George³, Manaf Alom¹, Zachary Kozel¹, Oksana Yaskiv¹, Simon J. Hall¹, Michael J. Schwartz¹, Manish A. Vira¹, Lee Richstone¹, Louis R. Kavoussi¹

1. Department of Urology, Smith Institute for Urology, Northwell Health, New Hyde Park, NY

2. Department of Urology, University of Illinois at Chicago, Chicago, IL

3. Department of Urology, University of Michigan, Ann Arbor, MI

Corresponding Author:

Paras H. Shah, MD
Smith Institute for Urology
405 Lakeville Road, Suite M41
New Hyde Park, NY 11040
Tel: 516-734-8500
Fax: 516-734-8619
E-Mail: pshah1023@gmail.com

Figures: 2

Tables: 3

Supplemental Tables: 2

Abstract Word Count: 237

Manuscript Word Count: 2499

Running Head: Pathologic T3a Upstaging after Partial and Radical Nephrectomy

Key Words: Partial nephrectomy, Radical nephrectomy, Extirpative surgery, Renal cell carcinoma, Kidney cancer

Funding: There were no external sources of funding

Download English Version:

<https://daneshyari.com/en/article/5686921>

Download Persian Version:

<https://daneshyari.com/article/5686921>

[Daneshyari.com](https://daneshyari.com)