Author's Accepted Manuscript

Determining when to stop PSA monitoring after radical prostatectomy: the role of ultrasensitive PSA

Kazuhiro Matsumoto , Akari Komatsuda , Yoshinori Yanai , Naoya Niwa , Takeo Kosaka , Ryuichi Mizuno , Eiji Kikuchi , Akira Miyajima , Mototsugu Oya



PII: S0022-5347(16)31195-8 DOI: 10.1016/j.juro.2016.08.098

Reference: JURO 13979

To appear in: The Journal of Urology

Accepted Date: 24 August 2016

Please cite this article as: Matsumoto K, Komatsuda A, Yanai Y, Niwa N, Kosaka T, Mizuno R, Kikuchi E, Miyajima A, Oya M, Determining when to stop PSA monitoring after radical prostatectomy: the role of ultrasensitive PSA, *The Journal of Urology*® (2016), doi: 10.1016/j.juro.2016.08.098.

DISCLAIMER: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our subscribers we are providing this early version of the article. The paper will be copy edited and typeset, and proof will be reviewed before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to The Journal pertain.

Embargo Policy

All article content is under embargo until uncorrected proof of the article becomes available online.

We will provide journalists and editors with full-text copies of the articles in question prior to the embargo date so that stories can be adequately researched and written. The standard embargo time is 12:01 AM ET on that date. Questions regarding embargo should be directed to jumedia@elsevier.com.

ACCEPTED MANUSCRIPT

Determining when to stop PSA monitoring after radical prostatectomy: the role of ultrasensitive PSA

Kazuhiro Matsumoto, Akari Komatsuda, Yoshinori Yanai, Naoya Niwa, Takeo Kosaka, Ryuichi Mizuno, Eiji Kikuchi, Akira Miyajima, and Mototsugu Oya

Keio University School of Medicine, Department of Urology, Shinanomachi 35, Shinjuku-ku, Tokyo 160-8582, Japan

Correspondence and reprint requests: Kazuhiro Matsumoto, M.D.

Keio University School of Medicine, Department of Urology,

Shinanomachi 35, Shinjuku-ku, Tokyo 160-8582, Japan

Tel: +81-3-5363-3825, Fax: +81-3-3225-1985, E-mail: kazz_matsumoto@yahoo.co.jp

Running head: Determining when to stop PSA monitoring

Key words: radical prostatectomy, PSA monitoring, follow-up, biochemical recurrence

Download English Version:

https://daneshyari.com/en/article/5687463

Download Persian Version:

https://daneshyari.com/article/5687463

<u>Daneshyari.com</u>