

Accepted Manuscript

Title: Prostatic Ductal Adenocarcinoma - a Rare Entity with Radiological Dilemma Demanding Endoscopic Biopsy

Author: S S Yadav, Pushpendra Baghel, Vinay Tomar, Neeraj Agarwal, Devendra Dhakad

PII: S0090-4295(17)30722-7

DOI: <http://dx.doi.org/doi: 10.1016/j.urology.2017.07.003>

Reference: URL 20548

To appear in: *Urology*

Received date: 12-4-2017

Accepted date: 3-7-2017

Please cite this article as: S S Yadav, Pushpendra Baghel, Vinay Tomar, Neeraj Agarwal, Devendra Dhakad, Prostatic Ductal Adenocarcinoma - a Rare Entity with Radiological Dilemma Demanding Endoscopic Biopsy, *Urology* (2017), <http://dx.doi.org/doi: 10.1016/j.urology.2017.07.003>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**Prostatic ductal adenocarcinoma - a rare entity with radiological dilemma
demanding endoscopic biopsy**

S S Yadav¹, Pushpendra Baghel^{*}, Vinay Tomar¹, Neeraj Agarwal¹, Devendra Dhakad¹,

¹ Department Of Urology, SMS medical college and associated hospital, Jaipur

***Corresponding author:** Pushpendra Baghel, MCh resident, Department of Urology, SMS medical college and associated hospital, Jaipur

Email: pushp.seoni@gmail.com

Phone: +918871089565

Conflict of interest: None

Source of financial support: None

Running title: The cystourethroscopic appearance and biopsy can guide in making diagnosis of prostatic ductal adenocarcinoma

Abstract

This is a rare case of prostatic ductal adenocarcinoma (PDA). A 75 year old gentleman presented with complaints of lower urinary tract symptoms (LUTS), constipation and occasional hematuria. The clinical and radiological differentiation from other retroperitoneal soft cystic lesion was difficult and FNAC from papillary solid components of cyst was required to reach diagnosis. TRUS guided biopsy was performed in view of raised serum PSA (> 100ng/ml), But, the cyst got ruptured

Download English Version:

<https://daneshyari.com/en/article/5690373>

Download Persian Version:

<https://daneshyari.com/article/5690373>

[Daneshyari.com](https://daneshyari.com)