

## Accepted Manuscript

Title: Incidence and risk factors of caesarean section in preterm breech births: a population-based cohort study

Authors: Elsa Lorthé, Mathilde Quere, Loïc Sentilhes, Pierre Delorme, Gilles Kayem



PII: S0301-2115(17)30122-7  
DOI: <http://dx.doi.org/doi:10.1016/j.ejogrb.2017.03.019>  
Reference: EURO 9823

To appear in: *EURO*

Received date: 2-2-2017  
Accepted date: 8-3-2017

Please cite this article as: Lorthé Elsa, Quere Mathilde, Sentilhes Loïc, Delorme Pierre, Kayem Gilles. Incidence and risk factors of caesarean section in preterm breech births: a population-based cohort study. *European Journal of Obstetrics and Gynecology and Reproductive Biology* <http://dx.doi.org/10.1016/j.ejogrb.2017.03.019>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# **Incidence and risk factors of caesarean section in preterm breech births: a population-based cohort study**

**Elsa Lorthé,<sup>1,2</sup> RM, MSc, Mathilde Quere,<sup>1</sup> MSc, Loïc Sentilhes,<sup>3</sup> MD, PhD, Pierre Delorme,<sup>1,4</sup> MD, Gilles Kayem,<sup>1,2,5</sup> MD, PhD**

<sup>1</sup> Inserm Unité Mixte de Recherche 1153, Obstetrical, Perinatal and Pediatric Epidemiology Research Team (Epopé), Center for Epidemiology and Statistics Sorbonne Paris Cité, Département Hospitalo-Universitaire Risks in pregnancy, Paris Descartes University

<sup>2</sup> Sorbonne Universités, Université Pierre & Marie Curie, Institut de Formation Doctorale, Paris, France

<sup>3</sup> Department of Obstetrics and Gynecology, Bordeaux University Hospital, Bordeaux, France

<sup>4</sup> Department of Obstetrics and Gynecology, Cochin, Broca, Hôtel Dieu Hospital, Assistance Publique Hôpitaux de Paris (AP-HP), Paris, France

<sup>5</sup> Department of Obstetrics and Gynecology, Trousseau Hospital, AP-HP, Paris, France

**Corresponding author:** Elsa Lorthé, INSERM U1153, Bâtiment Recherche, Hôpital Tenon, 4 rue de la Chine, 75020 Paris, France. Phone +33 1 56 01 83 67, fax +33 1 56 01 71 88. Email: elsa.lorthé@gmail.com

**Disclosure of interest:** The authors have no conflict of interest to report in relation with this article.

**Word count:** Abstract=283 words, Text=2542 words

**Condensation:** Breech presentation is common in preterm infants and is associated with widespread use of cesarean delivery despite the absence of recommendations for mode of delivery.

## **Abstract**

**Objectives:** To describe the incidence of breech presentation at 22 to 34 weeks' gestation, estimate the incidence of cesarean section delivery by cause of prematurity, and assess the factors associated with caesarean delivery in preterm breech births with preterm labor or preterm premature rupture of membranes.

**Study design:** EPIPAGE 2 is a French national prospective population-based cohort study of preterm births that occurred in 546 maternity units in 2011. We estimated the overall incidence of breech presentation and the incidence of cesarean delivery by cause of prematurity. Among the 579 singletons with breech presentation born at 22 to 34 weeks in a context of spontaneous

Download English Version:

<https://daneshyari.com/en/article/5692795>

Download Persian Version:

<https://daneshyari.com/article/5692795>

[Daneshyari.com](https://daneshyari.com)