



Contents lists available at ScienceDirect

Gynecology and Minimally Invasive Therapy

journal homepage: www.e-gmit.com

Original article

The benefit of adenomyomectomy on fertility outcomes in women with rectovaginal endometriosis with coexisting adenomyosis

Yohei Kishi ^{a, b, *}, Maki Yabuta ^b^a Department of Obstetrics and Gynecology, Nara Medical University, Nara, Japan^b Department of Obstetrics and Gynecology, Takanohara Central Hospital, U-Kyo 1-3-3, Nara 631-0805, Japan

ARTICLE INFO

Article history:

Received 22 December 2015
 Received in revised form
 9 August 2016
 Accepted 11 August 2016
 Available online xxx

Keywords:

adenomyosis
 endometriosis
 pregnancy
 rectovaginal
 surgery

ABSTRACT

Study Objective: To evaluate the effect of removal of coexisting adenomyosis on fertility outcomes in women with rectovaginal endometriosis.

Design: A retrospective cohort study.

Setting: A general hospital.

Patients: A total of 190 women who underwent laparoscopic nodule excision surgery for rectovaginal endometriosis between April 2007 and December 2012.

Interventions: Surgical excision of the rectovaginal endometriosis and coexisting uterine adenomyosis. Statistical analysis for fertility outcomes.

Measurement and main results: A total of 119 women desired postoperative pregnancy. Coexisting adenomyosis was found in 21% of the women. The overall clinical pregnancy rate was 41.2%. The only determining factor associated with a successful pregnancy was “age at surgery”. Clinical pregnancy rates with or without adenomyosis were 36.0% and 42.6%, respectively. We found no significant difference in clinical pregnancy rates between the groups.

Conclusion: There is a possibility that surgical removal of coexisting adenomyosis positively effects fertility outcomes in women with rectovaginal endometriosis. However, it is also important to note that the age at surgery was a critical factor for successful pregnancy.

Copyright © 2016, The Asia-Pacific Association for Gynecologic Endoscopy and Minimally Invasive Therapy. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Deep rectovaginal endometriosis is one of the most severe forms of pelvic endometriosis, and its prevalence is estimated to be 5–10% of all pelvic endometriosis cases.¹ With the recent trend toward delayed childbearing, patients with endometriosis associated subfertility show a tendency of aging and advancing in severity. This brings a range of difficulties in treatments aiming at restoring fecundity. In severe cases, adenomyosis often coexists with endometriosis and localizes at the outer myometrium without aberrations of the subendometrial myometrium² (Figure 1). Recent development of image diagnostic tools like magnetic resonance imaging (MRI) has enabled preoperative diagnosis of adenomyosis.

Now, the main treatment options of endometriosis associated subfertility are assisted reproductive technology (ART) and surgery. A recent article showed a higher clinical pregnancy rate of the combined treatment of surgery and subsequent *in vitro* fertilization (IVF) than that of surgery alone or IVF alone in women with endometriosis associated infertility.³ Regarding the determinants of fertility outcomes after surgical treatments for rectovaginal endometriosis, coexisting adenomyosis is often pointed out.^{4,5} However, as a recent review points out, adenomyosis was not excised in these reports.⁶ Thus, the effect of the removal of coexisting adenomyosis on fertility outcomes is not fully evaluated. We have performed a nodule excision surgery avoiding segmental resection in the vast majority of cases, and have routinely removed coexisting adenomyosis in all cases that MRI showed adenomyosis.

We set the aim of this study on analyzing factors relating to fertility outcomes in the surgery of deep rectovaginal endometriosis, and on evaluating the effect of removal of coexisting adenomyosis.

Conflicts of interest: The authors declare no conflicts of interest.

* Corresponding author. Department of Obstetrics and Gynecology, Takanohara Central Hospital, U-Kyo 1-3-3, Nara 631-0805, Japan.

E-mail address: kishi@takanohara-ch.or.jp (Y. Kishi).

<http://dx.doi.org/10.1016/j.gmit.2016.08.002>

2213-3070/Copyright © 2016, The Asia-Pacific Association for Gynecologic Endoscopy and Minimally Invasive Therapy. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article in press as: Kishi Y, Yabuta M, The benefit of adenomyomectomy on fertility outcomes in women with rectovaginal endometriosis with coexisting adenomyosis, Gynecology and Minimally Invasive Therapy (2016), <http://dx.doi.org/10.1016/j.gmit.2016.08.002>



Figure 1. Adenomyosis coexisting with pelvic endometriosis. T2-weighted magnetic resonance image (sagittal section) of an adenomyosis coexisting with pelvic endometriosis. The adenomyosis localizes at the outer myometrium. The junctional zone is kept intact without aberrancy and the healthy muscular structures can be seen in between the adenomyosis and the junctional zone. This image was taken in a 32-year-old nulliparous woman.

Materials and methods

We retrospectively compiled data of 190 women with histologically confirmed rectal endometriosis between April 2007 and December 2012. All the patients underwent surgery in the Department of Obstetrics and Gynecology, Takanohara Central Hospital, Nara, Japan. Surgical data including postoperative complications were retrieved from an operative database. Pregnancy outcomes were collected from questionnaires and/or interviews with outpatients. Among 174 women who underwent uterus conservative surgery, 119 desired postoperative pregnancies, and 116 (116/119) answered the questionnaire. In this study, we analyzed the 119 women that had the wish to get pregnant. Three women lost to follow-up were treated as “did not conceive”. We defined “clinical pregnancy” as the presence of a fetal heart beat at 12 weeks of gestation. All coexisting adenomyosis cases were diagnosed by preoperative MRI, and all of these were confirmed histologically. The criteria used for the definition of adenomyosis on MRI were: (1) a myometrial mass with indistinct margins of primarily low intensity with all sequences; or (2) diffuse or local widening of the junctional zone on T2 weighted images (wider than 12 mm).^{7–9} Furthermore, we analyzed the factors relating to clinical pregnancy by using univariable and multivariable regression analyses. The employed variables in these analyses were: age at surgery, coexisting adenomyosis, weight of adenomyosis, coexisting endometrioma, Revised American Fertility Society scores, whether performing a “full thick resection” or not, and presence or not of bilateral tubal occlusions in a tubal patency test during the surgery. We defined “full thick resection” as occurrence of rectal perforations during the slicing; a partial thick resection means perforation did not occur. None of the individuals were treated preoperatively with hormonal medicines. Preoperative pain symptoms were treated with nonsteroidal antiinflammatory drugs. Postoperative hormonal drugs were prescribed in some cases who did not wish for a pregnancy. This study was approved by the Institutional Ethical Review Board of Takanohara Central Hospital

on 28 November 2013 (ID 2013003), and informed consent was obtained from each of the patients.

Statistical analysis

The χ^2 test was used for the comparison of groups with regard to categorical variables; the Fisher exact test was used in the case of small cell counts. Parametric and nonparametric continuous variables were compared using the Student *t* test and the Mann–Whitney test. A *p* value < 0.05 was considered statistically significant. Stepwise logistic regression analysis was used for the multivariable analysis of the factors relating to clinical pregnancy (IBM Statistics software; version 16; SPSS Inc., Chicago, IL, USA).

Surgical procedures

The patient was placed in the Trendelenburg's position and triple puncture laparoscopic surgery was performed. We used a potassium titanyl phosphate laser device for incisions, and a bipolar device for vaporization. To superficial endometriosis, excision or vaporization was performed, and a cystectomy of ovarian endometrioma was carefully performed. Vaporization of the cyst wall was sometimes selected in cases having a history of ovarian cystectomy from the view of preservation of ovarian function. Rectovaginal endometriosis often infiltrates into the posterior wall of the uterus and the anterior wall of the rectum; the former one is often recognized as a posterior wall adenomyosis. With the use of a potassium titanyl phosphate laser device, we carefully sliced off the endometriotic nodules from both the anterior wall of the rectum and the posterior wall of the uterus. In instances having a very large nodule, we initiated the procedure by dissociation of the rectovaginal endometriotic nodules from the uterus, keeping the nodules attached to the anterior rectal wall, in the same fashion as other authors.^{10–12} Afterwards, the nodule was sliced off from the anterior wall of the rectum. The defected rectal wall (perforated or not) is continuously stitched with 2-0 synthetic absorbable sutures. For coexisting adenomyosis, we performed an adenomyomectomy as completely as possible in a surgical method reported earlier.¹³ Defected spaces were carefully obliterated with continuous stitches of 2-0 synthetic absorbable sutures to close the residual myometrium. All the surgeons performed these procedures routinely.

Results

Between 2007 and 2012, we performed 191 surgeries for deep rectovaginal endometriosis. Among them, 190 (190/191: 99.5%) cases were managed by laparoscopic nodule excision surgery with the exception of one case of segmental bowel resection operated with assistance from colorectal surgeons. Of the 174 women who selected conservative surgery, 119 desired postoperative pregnancy including five unmarried women (Table 1).

Pregnancy outcomes are presented in Table 2. The median follow-up period was 24 months (range: 12–60 months). Among the 119 women who wished to conceive or were unmarried at the time of surgery, 54/119 (45.3%) became pregnant. There was no pregnancy among five unmarried women. Clinical pregnancy was found in 49/119 (41.2%) women. All the clinical pregnancy cases were conceived spontaneously or by IVF (intracytoplasmic sperm injection [ICSI]-IVF).

All 49 cases of clinical pregnancy were delivered successfully, and none of the women who underwent adenomyomectomy delivered with the use of an elective cesarean section. The median time of conception linked to clinical pregnancy was 13.7 months (range: 1–56 months). There were no obstetrical complications. Spontaneous pregnancy was found in 34/119 (28.6%) cases, and ART

Download English Version:

<https://daneshyari.com/en/article/5695980>

Download Persian Version:

<https://daneshyari.com/article/5695980>

[Daneshyari.com](https://daneshyari.com)