

Accepted Manuscript

What is the most accurate lymph node staging method for perihilar cholangiocarcinoma? Comparison of UICC/AJCC pN stage, number of metastatic lymph nodes, lymph-node ratio, and log odds of metastatic lymph nodes

Simone Conci, MD, Andrea Ruzzenente, MD, PhD, Marco Sandri, MD, Francesca Bertuzzo, MD, Tommaso Campagnaro, MD, PhD, Fabio Bagante, MD, Paola Capelli, MD, Mirko D'Onofrio, MD, Marco Piccino, MD, Adriano E. Dorna, MD, Corrado Pedrazzani, MD, Calogero Iacono, MD, Prof., Alfredo Guglielmi, MD

PII: S0748-7983(17)30034-3

DOI: [10.1016/j.ejso.2016.12.007](https://doi.org/10.1016/j.ejso.2016.12.007)


Reference: YEJSO 4554

To appear in: *European Journal of Surgical Oncology*

Received Date: 11 August 2016

Revised Date: 20 October 2016

Accepted Date: 15 December 2016



Please cite this article as: Conci S, Ruzzenente A, Sandri M, Bertuzzo F, Campagnaro T, Bagante F, Capelli P, D'Onofrio M, Piccino M, Dorna AE, Pedrazzani C, Iacono C, Guglielmi A, What is the most accurate lymph node staging method for perihilar cholangiocarcinoma? Comparison of UICC/AJCC pN stage, number of metastatic lymph nodes, lymph-node ratio, and log odds of metastatic lymph nodes, *European Journal of Surgical Oncology* (2017), doi: 10.1016/j.ejso.2016.12.007.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

What is the most accurate lymph node staging method for perihilar cholangiocarcinoma?

Comparison of UICC/AJCC pN stage, number of metastatic lymph nodes, lymph-node ratio, and log odds of metastatic lymph nodes

Simone Conci, MD^{1*}; Andrea Ruzzenente, MD, PhD^{1*}; Marco Sandri, MD²; Francesca Bertuzzo, MD¹; Tommaso Campagnaro, MD, PhD¹; Fabio Bagante, MD¹; Paola Capelli, MD³; Mirko D'Onofrio, MD⁴; Marco Piccino, MD¹; Adriano E. Dorna, MD¹; Corrado Pedrazzani, MD¹; Calogero Iacono, MD¹§ and Alfredo Guglielmi, MD¹§

* equally contributed to the manuscript

§ shared the last authorship

¹ Department of Surgery, Division of General and Hepatobiliary Surgery, G.B. Rossi University Hospital, University of Verona, Verona, Italy

² Department of Economics and Management, Data Methods and System Statistical Laboratory, University of Brescia, Brescia, Italy

³ Department of Pathology and Diagnostics, G.B. Rossi University Hospital, University of Verona, Verona, Italy

⁴ Department of Radiology, G.B. Rossi University Hospital, University of Verona, Verona, Italy

Competing interests: none.

Correspondence to:

Prof. Calogero Iacono

Department of Surgery, Division of General and Hepatobiliary Surgery,

University of Verona Medical School, G.B. Rossi University Hospital

Piazzale L.A. Scuro, 10, Verona 37134, Italy

Tel.: +39-045-8124412; Fax: +39-045-8027426; Email: calogero.iacono@univr.it

Running Head: Lymph node staging methods for PCC

Download English Version:

<https://daneshyari.com/en/article/5701085>

Download Persian Version:

<https://daneshyari.com/article/5701085>

[Daneshyari.com](https://daneshyari.com)