



Case report

Optical coherence tomography angiography of diffuse unilateral subacute neuroretinitis

Ananda Kalevar ^{a, b}, J. Michael Jumper ^{a, b, *}^a Department of Ophthalmology, California Pacific Medical Center, San Francisco, CA, United States^b West Coast Retina Medical Group, San Francisco, CA, United States

ARTICLE INFO

Article history:

Received 2 December 2016

Received in revised form

27 March 2017

Accepted 20 June 2017

Available online 22 June 2017

Keywords:

Diffuse unilateral subacute neuroretinitis

DUSN

Optical coherence tomography angiography

OCTA

Imaging

ABSTRACT

Purpose: Diffuse unilateral subacute neuroretinitis (DUSN) is often a challenging diagnosis to make. We present a DUSN case with its multimodal imaging to aid in the diagnosis, emphasizing the observations on optical coherence tomography angiography (OCTA).

Observations: The evolution of a DUSN case is presented. Fundus photography and OCTA aided in the identification of the nematode.

Conclusion and importance: DUSN is a difficult diagnosis to establish. We report the first case to our knowledge in which OCTA aided in the diagnosis of DUSN.

© 2017 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Diffuse unilateral subacute neuroretinitis (DUSN) was initially described by Gass¹ in 1978. This paper came fifteen years after he and his colleagues recognized a “unilateral retinal wipe-out syndrome” in which healthy young individuals developed: “(1) insidious, usually severe loss of peripheral and central vision; (2) vitritis; (3) diffuse and focal pigment epithelial derangement with relative sparing of the macula; (4) narrowing of the retinal vessels; (5) optic atrophy; (6) increased retinal circulation time; and (7) subnormal electroretinographic findings.” (Gass and Scelfo).^{1–4} Patients can present in the early stage⁵ when the predominant findings are of chorioretinitis or in the late stage⁶ with poor vision, retinal and optic atrophy.

Most DUSN patients are young, present with altered vision and are in the late stages of the disease. Various nematode etiologies have been proposed depending on the size, speed of the nematode and the geographic area in which it was contracted.^{2–4,7} Most commonly, Baylisascaris procyonis and Ancylostoma caninum are thought to be the culprit. Precise diagnosis of the nematode is limited due to lack of good serologic testing or

stool examinations and most reports are made based on clinical exam. Specifically, that *B. procyonis* is longer and thought to be greater than 2000 μm while *A. caninum* is less between 250 and 500 μm.^{2–4}

Use of a contact lens to thoroughly examine the retina or examining color fundus photos have been generally recommended to help locate the worm. Despite these efforts and a strong clinical suspicion for DUSN, visualization of the worm on examination is uncommon and seen in only 25–40% of cases.^{2–4} OCT or OCTA have not been previously described in the identification of the worm in cases of DUSN. Laser photocoagulation applied to an identified worm is desired to limit retinal damage. Oral anthelmintics can be used in addition to laser to treat systemically⁸ or if a worm is not identified and the suspicion for DUSN is high.

2. Case report

A 49-year-old man presented with significantly decreased vision in the right eye for 2 months. He was previously treated with lisinopril for hypertension and, at presentation to our clinic, was on 80 mg prednisone for intraocular inflammation. He was a tree-trimmer by occupation in Northern California. Review of

* Corresponding author. 1445 Bush St, San Francisco, CA 94109, United States.

E-mail address: jmichaeljumper@gmail.com (J.M. Jumper).

systems revealed he had experienced widespread urticaria 1 month prior to presentation. No family history of ocular problems.

Visual acuity was hand motion (HM) and 20/16 in his right and left eye, respectively. A relative afferent papillary defect was present in the right eye. Intraocular pressure was within normal limits. The anterior segment examination was unremarkable. A posterior segment examination of the right eye revealed moderate vitritis, temporal optic disc pallor with severely attenuated retinal vessels (Fig. 1). In the macula, a motile subretinal worm (2500 μm x 100 μm) was identified with active retinitis seen superiorly (Fig. 2). Subretinal fibrosis and widespread retinal pigment epithelium (RPE) atrophy was also present. Serial fundus photography revealed significant progression from his exam 1 month earlier (Fig. 3). Posterior segment examination of the left eye was unremarkable.

Spectral-domain optical coherence tomography (SD-OCT) of the right eye revealed diffuse atrophy of both the inner and outer retinal layers as well as the RPE (Fig. 4). Prominent internal limiting membrane (ILM) was present (Oréface's sign). Wide-field fundus autofluorescence of the right eye showed diffuse, alternating, speckled hyper- and hypo-autofluorescence predominantly in the posterior pole (Fig. 5). Wide-field fluorescein angiography of the right eye revealed transmission hyperfluorescence in the areas of atrophy corresponding with window defect (Fig. 6). OCT angiography (OCTA) was performed on this patient and fortuitously, it revealed what we believed to be an anomalous pattern in the superotemporal macula (Fig. 7). As seen from the color fundus photos performed a few moments later, OCTA in fact revealed the mobile worm. The caliber and configuration seen on OCTA is consistent with a nematode. The cross-section OCTA reveals the worm within the retina curled upon itself (Fig. 8). Once the worm had moved again, OCTA was re-performed and it was confirmed that only normal retinal vasculature was seen and the anomalous pattern had disappeared. Laser photocoagulation was applied to the worm along its entire body. A 4-week course of oral albendazole was started and oral prednisone was tapered.



Fig. 1. Wide-field color fundus photo of the right eye revealing diffuse retinal pigment epithelium atrophy, fibrosis, attenuated vessels and optic disc pallor. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)




Fig. 2. Color fundus photo of the right macula revealing a worm. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)




Fig. 3. Color fundus photo of the right eye 1 month prior with what appears to be widespread retinal whitening. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

3. Discussion

Diffuse unilateral subacute neuroretinitis is an uncommon diagnosis. The prevalence of DUSN may actually be higher than reported due to the challenging course of the disease and the difficulty identifying the nematode. For example, there is a growing belief that many patients previously diagnosed with unilateral retinitis pigmentosa are in fact cases of undiagnosed DUSN.

Gass who initially described DUSN felt that high quality color fundus photos was the best way to identify nematodes. Our case proves that fundus photos still remain a valuable method of identification. However, with the worm not found in over 50% of suspected DUSN cases, there is still significant room for other techniques. To the best of our knowledge, this is the first report of OCTA aiding in the diagnosis of DUSN. Our case demonstrates

Download English Version:

<https://daneshyari.com/en/article/5703357>

Download Persian Version:

<https://daneshyari.com/article/5703357>

[Daneshyari.com](https://daneshyari.com)