Accepted Manuscript



Corneal endothelial cells activate innate and acquired arm of anti-viral responses after cytomegalovirus infection

Dai Miyazaki, Ryu Uotani, Michiko Inoue, Tomoko Haruki, Yumiko Shimizu, Keiko Yakura, Satoru Yamagami, Tatsuo Suzutani, Mayumi Hosogai, Hiroki Isomura, Yoshitsugu Inoue

PII: S0014-4835(17)30004-0

DOI: 10.1016/j.exer.2017.06.017

Reference: YEXER 7157

To appear in: *Experimental Eye Research*

Received Date: 11 January 2017

Revised Date: 21 June 2017

Accepted Date: 21 June 2017

Please cite this article as: Miyazaki, D., Uotani, R., Inoue, M., Haruki, T., Shimizu, Y., Yakura, K., Yamagami, S., Suzutani, T., Hosogai, M., Isomura, H., Inoue, Y., Corneal endothelial cells activate innate and acquired arm of anti-viral responses after cytomegalovirus infection, *Experimental Eye Research* (2017), doi: 10.1016/j.exer.2017.06.017.

Corneal endothelial cells activate innate and acquired arm of anti-viral responses after cytomegalovirus infection

Dai Miyazaki[1], Ryu Uotani[1], Michiko Inoue[1], Tomoko Haruki[1], Yumiko Shimizu[1], Keiko Yakura[1], Satoru Yamagami[2], Tatsuo Suzutani[3], Mayumi Hosogai[4], Hiroki Isomura[5], Yoshitsugu Inoue[1]

[1]Division of Ophthalmology and Visual Science, Faculty of Medicine, Tottori University, Tottori, Japan

[2]Department of Ophthalmology, Nihon University School of Medicine, Tokyo, Japan

[3]Department of Microbiology, Fukushima Medical University School of Medicine, Fukushima, Japan

[4]Department of Ophthalmology, Gunma University Graduate School of Medicine, Gunma, Japan

[5]Department of Virology, University of Toyama, Toyama, Japan

Abbreviation: cytomegalovirus: CMV, human corneal endothelial cell: HCEn cell, postinfection: PI, enzyme-linked immunosorbent assay: ELISA, indoleamine