Accepted Manuscript

The influence of early degenerative changes on the vulnerability of articular cartilage to impact-induced injury

Joshua Workman, Ashvin Thambyah, Neil Broom

PII: S0268-0033(17)30003-7

DOI: doi: 10.1016/j.clinbiomech.2017.01.003

Reference: JCLB 4264

To appear in: Clinical Biomechanics

Received date: 23 March 2016 Accepted date: 3 January 2017



Please cite this article as: Joshua Workman, Ashvin Thambyah, Neil Broom , The influence of early degenerative changes on the vulnerability of articular cartilage to impact-induced injury. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Jclb(2017), doi: 10.1016/j.clinbiomech.2017.01.003

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

The influence of early degenerative changes on the vulnerability of articular cartilage to impact-induced injury

Joshua Workman^a, Ashvin Thambyah^{a*}, Neil Broom^a

- a) University of Auckland, 2-6 Park Ave, Grafton, Auckland, 1023, New Zealand
- * = Corresponding author: Ashvin.thambyah@aucklanduni.ac.nz

All authors listed were fully involved in the study and preparation of the manuscript. The material contained within this manuscript has not been submitted or published elsewhere.

The authors declare no conflict of interest.

Word count:

Abstract: 249 Main text: 4104

Abstract:

Background

Recently, the structural changes in a bovine model of early degeneration were validated by our research group to be analogous to that in early human osteoarthritis. The hypothesis of this study was that the structural changes associated with increasing levels of degeneration would lead to higher levels of tissue damage in response to impact induced injury.

Methods

A total of forty bovine patellae were obtained for this study. Cartilage-on-bone samples were extracted from the distal lateral quarter, a region known to be affected by varying levels of

Download English Version:

https://daneshyari.com/en/article/5706902

Download Persian Version:

https://daneshyari.com/article/5706902

<u>Daneshyari.com</u>