

Accepted Manuscript

Title: Postural Control Strategies Differ in Normal Weight and Overweight Infants

Authors: Danae Dinkel, Kailey Snyder, Victoria Molfese, Anastasia Kyvelidou



PII: S0966-6362(17)30140-6
DOI: <http://dx.doi.org/doi:10.1016/j.gaitpost.2017.04.017>
Reference: GAIPOS 5390

To appear in: *Gait & Posture*

Received date: 14-12-2016
Revised date: 9-3-2017
Accepted date: 12-4-2017

Please cite this article as: Dinkel Danae, Snyder Kailey, Molfese Victoria, Kyvelidou Anastasia. Postural Control Strategies Differ in Normal Weight and Overweight Infants. *Gait and Posture* <http://dx.doi.org/10.1016/j.gaitpost.2017.04.017>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title: Postural Control Strategies Differ in Normal Weight and Overweight Infants**Running title:** Infant Postural Control & Weight

Danae Dinkel^a, Kailey Snyder^a, Victoria Molfese^b, Anastasia Kyvelidou^c

^aSchool of Health and Kinesiology
College of Education
University of Nebraska at Omaha
6001 Dodge Street
Omaha, NE 68182
United States of America

^bDepartment of Child, Youth, and Family Studies
College of Education and Human Sciences
University of Nebraska at Lincoln
1400 R St
Lincoln, NE 68588
United States of America

^c Department of Biomechanics
College of Education
6160 University Drive
Omaha NE 68182
United States of America

Contact Information:

Danae Dinkel, Ph.D. (**Corresponding author**)
dmdinkel@unomaha.edu

Kailey Snyder
kesnyder@unomaha.edu

Victoria Molfese
vmolfese2@unl.edu

Anastasia Kyvelidou
akyvelidou@unomaha.edu

Highlights

- Normal weight and overweight infants learned to sit at similar ages.
- Postural control strategies differed between the two groups at sitting onset.
- Normal weight infants had greater postural sway at one-month post onset of sitting.

Download English Version:

<https://daneshyari.com/en/article/5707878>

Download Persian Version:

<https://daneshyari.com/article/5707878>

[Daneshyari.com](https://daneshyari.com)