Accepted Manuscript

Title: Trunk biomechanics during hemiplegic gait after stroke: a systematic review

Authors: Van Criekinge Tamaya, Saeys Wim, Hallemans Ann, Velghe Silke, Viskens Pieter-Jan, Vereeck Luc, De Hertogh Willem, Truijen Steven



DOI: http://dx.doi.org/doi:10.1016/j.gaitpost.2017.03.004

Reference: GAIPOS 5338

To appear in: Gait & Posture

Received date: 30-4-2016 Revised date: 27-2-2017 Accepted date: 2-3-2017

Please cite this article as: Tamaya Van Criekinge, Wim Saeys, Ann Hallemans, Silke Velghe, Pieter-Jan Viskens, Luc Vereeck, Willem De Hertogh, Steven Truijen.Trunk biomechanics during hemiplegic gait after stroke: a systematic review. *Gait and Posture* http://dx.doi.org/10.1016/j.gaitpost.2017.03.004

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

Trunk biomechanics during hemiplegic gait after stroke: a systematic review

Van Criekinge Tamaya, MSc^a (corresponding author)

tamaya.vancriekinge@uantwerpen.be, +32472540078

Universiteit Antwerpen, Department of Rehabilitation Sciences and Physiotherapy,

Universiteitsplein 1, 2610 Antwerp.

Saeys Wim, PhDa

Hallemans Ann, PhDa

Velghe Silke, MSc

Viskens Pieter-Jan, MSc

Vereeck Luc, PhDa

De Hertogh Willem, PhDa

Truijen Steven, PhDa

^aDepartment of Rehabilitation Sciences and Physiotherapy, Faculty of Medicine and Health Sciences, University of Antwerp, Universiteitsplein 1, 2610 Antwerp, Antwerp, Belgium

e-mail: willem.saeys@uantwerpen.be ; luc.vereeck@uantwerpen.be ; steven.truijen@uantwerpen.be ; steven.truijen@uantwerpen

Highlights

- Stroke patients walk with less dissociation between thorax and pelvis
- After stroke less stability and symmetry is observed in thorax and pelvis
- Stroke patients show increased mediolateral and anteroposterior trunk movements
- Increasing walking speed normalizes thoracic and pelvic movements

Download English Version:

https://daneshyari.com/en/article/5707892

Download Persian Version:

https://daneshyari.com/article/5707892

<u>Daneshyari.com</u>