



Clinical case

Rhomboid nerve transfer to the suprascapular nerve for shoulder reanimation in brachial plexus palsy: A clinical report

Transfert du nerf du rhomboïde sur le nerf suprascapulaire pour la réanimation de l'épaule dans une paralysie du plexus brachial : un cas clinique

J.-N. Goubier*, F. Teboul

Institute of brachial plexus and nerve surgery, 92, boulevard de Courcelles, 75017 Paris, France

Received 16 April 2016; received in revised form 8 June 2016; accepted 3 July 2016

Available online 23 September 2016

Abstract

Recovery of shoulder function is a real challenge in cases of partial brachial plexus palsy. Currently, in C5–C6 root injuries, transfer of the long head of the triceps brachii branch is done to revive the deltoid muscle. Spinal accessory nerve transfer is typically used for reanimation of the suprascapular nerve. We propose an alternative technique in which the nerve of the rhomboid muscles is transferred to the suprascapular nerve. A 33-year-old male patient with a C5–C6 brachial plexus injury with shoulder and elbow flexion palsy underwent surgery 7 months after the injury. The rhomboid nerve was transferred to the suprascapular nerve and the long head of the triceps brachii branch to the axillary nerve for shoulder reanimation. A double transfer of fascicles was performed, from the ulnar and median nerves to the biceps brachii branch and brachialis branch, respectively, for elbow flexion. At 14 months' follow-up, elbow flexion was rated M4. Shoulder elevation was 85 degrees and rated M4, and external rotation was 80 degrees and rated M4. After performing a cadaver study showing that transfer of the rhomboid nerve to the suprascapular nerve is technically possible, here we report and discuss the clinical outcomes of this new transfer technique.

© 2016 Published by Elsevier Masson SAS on behalf of SFCM.

Keywords: Rhomboid nerve; Suprascapular nerve; Brachial plexus palsy; Nerve transfer

Résumé

La réanimation de l'épaule dans le cadre des paralysies du plexus brachial demeure un vrai challenge. Actuellement, dans le cadre d'une lésion de la racine C5, un transfert du nerf du chef long du triceps brachial sur le nerf axillaire est proposé pour la réinnervation du deltoïde. Concernant la réanimation du nerf suprascapulaire, le nerf spinal accessoire est actuellement le plus utilisé. Nous proposons une alternative à ce transfert utilisant le transfert du nerf des rhomboïdes sur le nerf suprascapulaire. Un homme de 33 ans, présentant une rupture des racines C5C6 était opéré à 7 mois de l'accident. Concernant la réanimation de l'épaule, un transfert du nerf des rhomboïdes sur le nerf suprascapulaire était pratiqué ainsi qu'un transfert du nerf du chef long du triceps brachial sur la branche antérieure du nerf axillaire. Concernant la flexion du coude, un double transfert de fascicules du nerf ulnaire et du nerf médian sur les nerfs du biceps brachial et du brachial était réalisé. Au suivi de 14 mois, l'élévation de l'épaule était de 85° et la rotation latérale de 80° cotées M4. La flexion du coude était complète et cotée M4. Après une étude anatomique prouvant la faisabilité de ce transfert, nous rapportons et discutons les résultats cliniques de ce nouveau transfert.

© 2016 Publié par Elsevier Masson SAS au nom de SFCM.

Mots clés : Nerf du rhomboïde ; Nerf suprascapulaire ; Paralysie du plexus brachial ; Transfert nerveux

* Corresponding author.

E-mail address: jngoubier@gmail.com (J.N. Goubier).

1. Introduction

Shoulder palsy in brachial plexus injuries of the C5 and C6 roots is a real challenge for nerve surgeons. Transfers from the radial nerve branches to the axillary nerve are generally used for anterior deltoid muscle reanimation [1,2]. As for the suprascapular nerve (SSN), the spinal accessory nerve (SAN) is usually transferred for reanimation of the supraspinatus and infraspinatus muscles [3,4]. We propose an alternative technique for reanimation of the SSN using the nerve to the rhomboid muscles, in order to keep the trapezius muscle intact, or potentially to use the SAN for another target.

A cadaver study was performed initially to prove the feasibility of this transfer (Fig. 1). The results showed that tensionless suture of the rhomboid nerve to the SSN was possible in all shoulders, with the diameter of the two nerves being macroscopically compatible (average diameter: 2.9 and 3 mm, respectively) [5]. We now present the results of the transfer of the rhomboid nerve to the SSN in a clinical case.

2. Clinical report

A 33-year-old male patient had partial palsy of his right brachial plexus involving the C5 and C6 roots following a motorcycle accident. Shoulder and elbow flexion were paralyzed. The anterior serratus and rhomboid muscles were clinically present. Electromyography confirmed the clinical examination and MRI excluded root avulsions.

Surgical exploration was deemed appropriate. Time before surgery was 7 months. Transfer from one fascicle of the ulnar




Fig. 1. Transfer of the rhomboid nerve to the suprascapular nerve: the levator scapulae is released from the medial border of the scapula to expose the dorsal scapular nerve. The nerve to the rhomboid muscles is released until it reaches the upper edge of the rhomboid muscles, divided, and then turned toward the suprascapular nerve in the supraspinatus fossa (from Goubier and Teboul [5]).

and median nerve to the biceps and brachial branches was performed. For the shoulder palsy, transfer from the long head of the triceps branch (LHTB) to the anterior branch of the axillary nerve for the anterior deltoid muscle was carried out. A transfer from the rhomboid nerve to the SSN was also performed.

The patient was positioned in lateral decubitus. A posterior approach along the spine of the scapula was performed. The trapezius was detached close to its scapular insertion in order to access the supraspinatus fossa. Then, the deep aspect of the supraspinatus muscle was reflected in order to access the suprascapular notch. At this level, the SSN was located, and freed by sectioning the transverse ligament of scapula to obtain the maximal length possible so as to make subsequent suturing easier.

The levator scapulae muscle was detached from the medial border of the scapula, and then retracted to expose the deep aspect of the dorsal scapular nerve. Once exposed, the branches to the levator scapulae muscle and rhomboid muscle were isolated. Electrical stimulation of the rhomboid nerve, starting with low intensity (0.02 mA) was performed to confirm normal innervation. The branch for the rhomboid muscle was freed as far as possible in the muscle and in the dorsal scapular nerve in order to obtain the maximum possible length. The branch was then cut close to the muscle and brought into contact with the SSN (Fig. 2). Then, the rhomboid and SSN were sutured under microscopy, with three separate 11-0 nylon sutures supplemented with fibrin glue. The trapezius muscle was reinserted to the spine of scapula. The skin was closed without a suction drain. The upper limb was immobilized in a splint for three weeks, after which rehabilitation was started with passive motion of the upper limb. Once the first contractions occurred, electrical stimulation of re-innervated muscles was performed.

At 7 months' follow-up, elbow flexion range-of-motion was complete and strength was rated M4 (according to Medical Research Council scoring). At 14 months' follow-up, active shoulder forward flexion was 85 degrees and rated M4. Active external rotation of the shoulder (measured from full internal




Fig. 2. Transfer of the rhomboid nerve to the suprascapular nerve with a posterior approach (right shoulder). This transfer can be performed without any tension. With this approach, the nerve is sutured close to the supraspinatus muscle, encouraging faster recovery.

Download English Version:

<https://daneshyari.com/en/article/5708369>

Download Persian Version:

<https://daneshyari.com/article/5708369>

[Daneshyari.com](https://daneshyari.com)