Accepted Manuscript

Can surgeons reduce the risk for dislocation following primary total hip arthroplasty performed using the posterolateral approach?

Kurt G. Seagrave, Anders Troelsen, Bjørn Gliese, Henrik Husted, Thomas Kallemose, Kirill Gromov

PII: S0883-5403(17)30388-1

DOI: 10.1016/j.arth.2017.04.047

Reference: YARTH 55849

To appear in: The Journal of Arthroplasty

Received Date: 12 January 2017
Revised Date: 19 March 2017
Accepted Date: 24 April 2017

Please cite this article as: Seagrave KG, Troelsen A, Gliese B, Husted H, Kallemose T, Gromov K, Can surgeons reduce the risk for dislocation following primary total hip arthroplasty performed using the posterolateral approach?, *The Journal of Arthroplasty* (2017), doi: 10.1016/j.arth.2017.04.047.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT Can surgeons reduce the risk for dislocation following primary total hip arthroplasty performed using the posterolateral approach?

Kurt G Seagrave ¹
Anders Troelsen ¹
Bjørn Gliese ¹
Henrik Husted ¹
Thomas Kallemose 1, 2
Kirill Gromov ¹
¹ Department of Orthopaedic Surgery, Copenhagen University Hospital Hvidovre. Copenhagen, Denmark
² Clinical Research Centre, Copenhagen University Hospital Hvidovre. Copenhagen, Denmark
Corresponding author: Kirill Gromov
Email: kirgromov@gmail.com Postal address: Department of Orthopaedic Surgery. Hvidovre Hospital. Kettegård Allé 30.
2650. Copenhagen, Denmark

Download English Version:

https://daneshyari.com/en/article/5708587

Download Persian Version:

https://daneshyari.com/article/5708587

<u>Daneshyari.com</u>