Accepted Manuscript

Title: Maximum Pain on Visual Analog Scales in Spinal Disorders

Author: Caleb J. Behrend, Etienne M. Schönbach, Alexander R. Vaccaro, Ellen Coyne, Mark L. Prasarn, Glenn R. Rechtine

PII: S1529-9430(16)31111-1
DOI: http://dx.doi.org/doi: 10.1016/j.spinee.2016.11.017
Reference: SPINEE 57215

To appear in: *The Spine Journal*

Received date: 21-8-2016
Revised date: 28-10-2016
Accepted date: 28-11-2016

Please cite this article as: Caleb J. Behrend, Etienne M. Schönbach, Alexander R. Vaccaro, Ellen Coyne, Mark L. Prasarn, Glenn R. Rechtine, Maximum Pain on Visual Analog Scales in Spinal Disorders, *The Spine Journal* (2016), http://dx.doi.org/doi: 10.1016/j.spinee.2016.11.017.

Title Page

Maximum pain on visual analog scales in spinal disorders.

Caleb J. Behrend, MD[1]

Etienne M. Schönbach, MD[2]

Alexander R. Vaccaro, MD, PhD[3]

Ellen Coyne, MS[4]

Mark L. Prasarn, MD[5]

Glenn R. Rechtine, MD[6]

1 Carilion Clinic Orthopaedics, Virginia Tech, Roanoke, VA, USA

2 Wilmer Eye Institute, Johns Hopkins University School of Medicine, Baltimore, MD, USA

3 Rothman Institute, Thomas Jefferson University, Philadelphia, PA, USA

4 Independent Statistical Support, Rochester, NY, USA

5 Department of Orthopaedics and Rehabilitation, University of Texas, Houston, TX, USA

6 Department of Orthopaedics, Veterans Affairs Medical Center, Asheville, NC, USA

Corresponding Author:

Etienne Schönbach, MD,

Wilmer Eye Institute, Johns Hopkins University School of Medicine, Woods 259,

600 N. Wolfe Street, Maryland 21287, USA

1 Abstract

Background context: Determining pain intensity is largely dependent on the patient's report.