



Review

Porocarcinoma; presentation and management, a meta-analysis of 453 cases



Abdulwahid M. Salih ^a, F.H. Kakamad ^{b, c, *}, Hiwa O. Baba ^d, Rawezh Q. Salih ^{c, e},
M.R. Hawbash ^{c, f}, Shvan H. Mohammed ^{c, e}, Snur Othman ^d, Yadgar A. Saeed ^d,
Imad J. Habibullah ^d, Aso S. Muhiyaldeen ^d, Rebaz O. Nawroly ^d, Zuhair D. Hammood ^d,
Nawshirwan H. Abdulkarim ^d

^a Faculty of Medical Sciences, School of Medicine, Department Surgery, University of Sulaimani, François Mitterrand Street, Sulaimani, Kurdistan, Iraq

^b Faculty of Medical Sciences, School of Medicine, Department Cardiothoracic and Vascular Surgery, University of Sulaimani, François Mitterrand Street, Sulaimani, Kurdistan, Iraq

^c Bioscience Center, Hamdi Street, Sulaimani, Kurdistan, Iraq

^d Sulaimani Teaching Hospital, Sulaimani, Kurdistan, Iraq

^e Faculty of Science & Science Education, School of Science, Biology Department, University of Sulaimani, Raperin Street, Sulaimani, Kurdistan, Iraq

^f Faculty of Science & Science Education, School of Science Education, Biology Department, University of Sulaimani, François Mitterrand Street, Sulaimani, Kurdistan, Iraq

H I G H L I G H T S

- Porocarcinoma is a rare sort of skin cancer developing from sweat glands.
- Because of rarity of the problem, there is controversy regarding both presentation and management.
- The aim of the current meta-analysis is to summarize all data regarding the disease.

A R T I C L E I N F O

Article history:

Received 22 March 2017

Received in revised form

18 June 2017

Accepted 18 June 2017

Keywords:

Porocarcinoma

Presentation

Management

Aggressive

Carcinoma

A B S T R A C T

Background: porocarcinoma is a rare sort of skin cancer developing from sweat glands. Its clinical course and management are not well understood.

Objectives: the current meta-analysis is to address the presentation and management of porocarcinoma. **Data sources:** Web of Science, PubMed, MEDLINE on OVID and Google scholar were searched for English-language studies published before December 1, 2016.

Results: The review of literature revealed 453 cases. From which 222 (49%) cases were male and female were 231 (51%). The mean age was 67.57 years. The mean duration of presentation was 5.57 years ranging from 4 days to 60 years of age. The most common site of affection is the head and neck (39.9%) followed by lower extremity (33.9%). Mass and nodule are the most common modes of presentation. Metastasis occurred at presentation in 110 (31%) cases. The most common organ to which porocarcinoma metastasizes is the nearby lymph node (57.7%).

Conclusions: Porocarcinoma is an aggressive skin cancer. Surgery is the main modality of treatment.

Systematic review registration number: reviewregistry233.

© 2017 The Author(s). Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author. Faculty of Medical Sciences, School of Medicine, Department Cardiothoracic and Vascular Surgery, University of Sulaimani, François Mitterrand Street, Sulaimani, Kurdistan, Iraq.

E-mail address: fahmi.hussein@univsul.edu.iq (F.H. Kakamad).

Contents

1. Introduction	75
2. Methodology	75
3. Result	75
4. Discussion	75
Ethical approval	77
Sources of funding	77
Author contribution	77
Conflicts of interest	77
Research registration unique identifying number (UIN)	77
Guarantor	77
References	77
Further reading	79

1. Introduction

Porocarcinoma is a rare sort of skin cancer developing from sweat glands, specifically, it is a malignancy of the eccrine sweat glands [1]. Its etiology is not well understood. However, some studies have showed that the tumor developed from a pre-existing eccrine poroma. From others perspective, chronic light exposure, exposure to chemical agents and immunosuppression could be the predisposing factors for de novo eccrine porocarcinoma [2]. Porocarcinoma is a dangerous disease because of high rate of recurrence after resection and metastasis to vital organs [3,4]. Because of rarity of the problem, there is controversy regarding both presentation and management [5,6]. Liyod et al. revealed features like a rapid increase in size, foul odor and fungating fleshy appearance while Kurashige and colleagues who collects 8 cases revealed that all cases of porocarcinoma presented with erosive reddish nodules [6,7].

Controversy regarding treatment is also present [8]. Kurashige and colleagues who present 8 cases of porocarcinoma summarized that surgical resection should be done in all cases, as adjuvant therapy in form of local irradiation, while Fujimura et al. preferred surgical resection of the mass and giving adjuvant therapy only in special situation [7]. The aim of the current meta-analysis is to summarize all data regarding the presentation and management of cases with porocarcinoma which was done in line with PRISMA guide line [9].

2. Methodology

Information sources and Search: Web of Science, PubMed, MEDLINE on OVID and Google scholar were searched for English-language studies published before December 1, 2016. The search key words were porocarcinoma, carcinoma of sweat gland and tumor of sweat gland. The data collection was supplemented by references cited by included articles.

Eligibility criteria: For a study to be included in this meta-analysis, it should address at least one specific aim (clinical course, management, or both) of porocarcinoma (Fig. 1).

Data collection process: Data were taken directly from the articles by 4 authors independently. Investigators have not been contacted to obtain and confirm the data.

Data items: Several data were extracted and pooled from all included articles. Those articles were socio-demographic characteristics of the patients, sample size, presentations, duration of presentation, method of diagnosis, options for managements, recurrence rate and complications.

Summary measures and Synthesis of results: Extracted data were calculated and re-analyzed totally. According to the variables, they are showed as percentage, mean values, and ranges of variation and percentages. The search in the medical literature revealed 104 studies which included 453 patients.

3. Result

The result reveals 453 patients. From which 222 (49%) case were male and female were 231 (51%). The mean age was 67.57 years ranging from 6 months to 97 years of age. The mean duration of presentation was 5.57 years ranging from 4 days to 60 years. The most common sites of affection are the head and neck (39.9%) followed by lower extremity (33.9%). Table 1 shows the distribution of the patients according to the site of affection. Mass and nodule are the most common modes of presentation. Table 2 shows presentation of the porocarcinoma with their frequency of appearance. Metastasis occurred at presentation in 110 (31%) cases. The most common organs to which porocarcinoma metastasize are the nearby lymph nodes (57.7%), followed by lung (12.8%). Table 3 shows the organs of metastasis.

4. Discussion

Based on their findings, some authors stated that porocarcinoma is more common among male patients [7,26]. The current metadata showed that porocarcinoma is evenly distributed according to the gender, both male and female are equally affected, (male 49% is nearly the same female 51%). The mean age of presentation is seventh and eighth decades of life (mean age 67.57 years, ranging from 7 months to 97 years) [34]. Most of the patients present with mass without symptoms (poroma) long time before converting to malignant type (porocarcinoma) [8,62]. According to this study, the mean duration of presentation was 5.73 years. Robson and colleagues reported a case of porocarcinoma presented with history of 60 years of benign stable mass [30]. This finding may suggest that aggressive management of benign subcutaneous mass is advised to prevent malignant conversion especially when the diagnosis is not confirmed to be benign. The literature showed that most of the cases of porocarcinoma present with either nodule or mass [31]. The current meta-data also confirmed this finding and about 71.2% cases presented with either nodule or mass. Yamamoto and associates reported that majority of porocarcinoma lesions appear in the upper extremity (80%) while Shiohara and colleagues revealed the reverse of this finding. In their 12 case series, they reported that in 9 (75%) cases of them, the primary site was the

Download English Version:

<https://daneshyari.com/en/article/5722880>

Download Persian Version:

<https://daneshyari.com/article/5722880>

[Daneshyari.com](https://daneshyari.com)