



Case report

A case report of an unusual cause of postoperative leg pain after posterior lumbar fusion



R. Daureeawoo Foundation Year 2 Doctor ^{a, b, *}, S. Baliga Spinal Fellow ^a,
W. Mohahmed Vascular fellow ^c, E.B. Ahmed Consultant Spinal Surgeon ^a

^a Spinal Surgery Unit, Department of Trauma and Orthopaedics, University Hospitals of North Midlands NHS Trust, Newcastle Road, Stoke-on-Trent, ST4 6QG, United Kingdom

^b Accident & Emergency Department, University Hospitals of North Midlands NHS Trust, United Kingdom

^c Department of General Surgery, University Hospitals of North Midlands NHS Trust, United Kingdom

HIGHLIGHTS

- Recognition of post-operative complication in minimally invasive spine surgery.
- Importance of urgency in investigating in post-operative spine patients.
- Misplacement of a screw in spine surgery can have serious and potentially fatal outcomes in patients.

ARTICLE INFO

Article history:

Received 22 March 2017

Received in revised form

19 May 2017

Accepted 20 May 2017

Keywords:

Post-operative complication

Case report

Haemorrhage

Pain

Spinal fusion

ABSTRACT

We report the case of a 36-year old man who underwent elective posterior lumbar spinal fusion after presenting with bilateral leg pain with associated back pain. He had undergone a L5/S1 discectomy in 2001, which provided good symptomatic relief. On this admission, he underwent elective L4-S1 posterolateral fusion and bilateral L4/5 and L5/S1 decompression. Intra-operatively a pedicle screw had to be re-inserted after fluoroscopy confirmed a lateral breach. The patient had no major postoperative complications until the sixth day when the patient re-presented with acute leg pain and weakness. Following a local multidisciplinary meeting (MDT) an MRI showed a large haematoma at the right psoas muscle. CT angiogram confirmed a bleeding lumbar segmental vessel at the site of the previous misplaced screw and an emergency fluoroscopic guided embolisation of the vessel was performed. The patient recovered well post operatively and was discharged back to the community 12 days later. Haemorrhage, as in this case a psoas haematoma should be considered as one of the differentials in patients with recurrent post-operative radiculopathy following posterior lumbar spine fusion. This should be suspected particularly if there has been a misplacement of one or more screws as this can cause injury to blood vessels and be fatal for the patient.

Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Post-operative complications are often experienced in a specialised area of surgery that requires specific care for the intervention performed. Lumbar artery haemorrhage is in most cases associated

with anticoagulant therapy [1] and injuries following lumbar spine surgery have rarely been described [2–4]. Although rare, we should be aware of the significant mortality associated with haemorrhage in minimally invasive spine surgery. When suspected, urgent intervention is recommended for best patient outcome.

2. Case presentation

A 36-year-old gentleman underwent elective L4-S1 posterolateral fusion and bilateral L4/5 L5/S1 decompression. Fourteen years previously, he had undergone a left sided L5/S1 discectomy, which at the time had given him excellent relief of his symptoms.

* Corresponding author. Spinal Surgery Unit, Department of Trauma and Orthopaedics, University Hospitals of North Midlands NHS Trust, Newcastle Road, Stoke-on-Trent, ST4 6QG, United Kingdom.

E-mail addresses: rd9922@my.bristol.ac.uk (R. Daureeawoo), santoshbaliga@doctors.org.uk (S. Baliga), Wafaa.mohammed@uhns.nhs.uk (W. Mohahmed), Elnasri.ahmed@uhns.nhs.uk (E.B. Ahmed).

Subsequently, he was referred to our unit for further investigation and management. He reported a 12-month history of bilateral leg pain with associated back pain. The leg pain was consistent with bilateral L5 radiculopathy. Standing radiographs of the lumbar spine showed reduced disc heights of the L4/5 and L5/S1 levels but were otherwise unremarkable. An MRI revealed a broad based central L4/5 disc prolapse resulting in canal stenosis and bilateral nerve root compression and well as a of bilateral L5 compression from foraminal narrowing secondary to loss of disc height and circumferential osteophytic bar (See Fig. 1).

Bilateral L5 nerve root blocks were performed which gave the patient three weeks of complete relief of his symptoms; after which his leg pain slowly returned to previous levels. Hence the patient was offered and consented to L4/5 decompression and discectomy, L5/S1 decompression with L4-S1 posterolateral fusion. The aim of this was to decompress all neural elements, in particular the L5 nerve roots.

The patient underwent the above elective procedure under general anaesthetic and neural monitoring. Pedicle tracks were established using free-hand technique, standard anatomic landmarks and two-person probing. After palpation of bone in all five walls, a marker was placed. Fluoroscopic images, AP and lateral were obtained to further confirm accurate tracks. This was then followed by a thorough decompression of both levels.

Pedicle screws were then inserted into the tracks, sequentially replacing the markers. All screws had excellent hold apart from the right L4 screw. Fluoroscopy confirmed that this screw had not followed the original track and had breached laterally. The screw was hence removed, the correct track retraced and the screw reinserted. Repeat fluoroscopy showed good position of all screws. The rest of the procedure went without complication and no significant bleeding was encountered at anytime.

Immediately after the operation, the patient complained of numbness in the right thigh (L2 distribution). This was accompanied by weakness of hip flexion, 4 out of 5 on the Medical Research Council Scale for Muscle Strength [5]. However these symptoms resolved over the course of 12 hours and was put down to the epidural analgesia that had been administered intra-operatively.

On the first post-operative day, his Haemoglobin (Hb) was 104 g/L, the neurology was normal and the patient was mobilised

with routine physiotherapy. Three days after surgery a check x-ray of his lumbar spine was performed and this showed good spinal alignment and satisfactory position of fixation. The patient was discharged home on the fifth post-operative day.

The patient returned to the Emergency Department 24 hours later, with pins and needles in the right thigh, pain in the right groin and reduced power in his right hip. His bladder and bowel functions were normal and the surgical wound site showed no signs of infection. Femoral nerve stretch was strongly positive on the right. Repeat radiographs of the lumbar spine were identical to those taken prior to discharge but an MRI scan revealed a massively enlarged right psoas muscle and the appearance suggested the presence of a large haematoma and secondary inflammation (See Fig. 2).

Although the patient was haemodynamically stable, blood investigations revealed a Hb of 84 g/L which fell to 72 g/L six hours later. After discussion with the radiologist on duty, we decided to proceed with a CT angiogram. This revealed a pseudoaneurysm arising from the right L4 lumbar artery, which was considered to not only be the source for the psoas haematoma but also to be actively bleeding.

An emergency fluoroscopic guided embolisation of the vessel under local anaesthetic was performed and there were no immediate procedural complications.

Over the following 48 hours, his groin pain improved as did the power in his right lower-limb. He was mobilised with physiotherapy and was discharged home several days later. On review in the clinic about 6 months later, he still complained of mild pain to his back and right leg. However, he had returned to work and has been having ongoing physiotherapy in the community. An MRI was arranged and the results were satisfactory (See Fig. 3). He was referred back to the physiotherapist to continue his recovery.

3. Discussion

Cadaveric studies have shown that the first four lumbar arteries originate directly from the aorta, while the fifth lumbar artery can branch from the iliolumbar artery or the median sacral artery. The



Fig. 1. Sagittal slice of MRI showing disc dessication at L4/5 and L5/S1 levels.



Fig. 2. Post-op MRI showing large haematoma in the right psoas.

Download English Version:

<https://daneshyari.com/en/article/5722920>

Download Persian Version:

<https://daneshyari.com/article/5722920>

[Daneshyari.com](https://daneshyari.com)