



# Full fat milk consumption protects against severe childhood obesity in Latinos☆☆☆

Amy L. Beck<sup>a</sup>, Melvin Heyman<sup>b</sup>, Cewin Chao<sup>c</sup>, Janet Wojcicki<sup>b,\*</sup>

<sup>a</sup> University of California San Francisco, 3333 California St. Suite 245, Box 0503, San Francisco, CA, 94118, United States

<sup>b</sup> University of California San Francisco, 550 16th Street 5th Floor, Mail Stop 0136, San Francisco, CA, 94143, United States

<sup>c</sup> University of California San Francisco, 505 Parnassus Ave., San Francisco, CA, 94143, United States

## ARTICLE INFO

### Article history:

Received 13 February 2017

Received in revised form 25 May 2017

Accepted 14 July 2017

Available online 23 July 2017

### Keywords:

Preschoolers

Dairy products

Latinos

Nutrition policy

Health disparities

## ABSTRACT

Consumption of non- or low-fat dairy products is recommended as a strategy to lower the risk of childhood obesity. However, recent evidence suggests that consumption of whole fat dairy products may, in fact, be protective against obesity. Our objective was to determine the association between milk fat consumption and severe obesity among three-year-old Latino children, a population with a disproportionate burden of obesity and severe obesity. 24-hour-dietary recalls were conducted to determine child intake in San Francisco based cohort recruited in 2006–7. Mother-child dyads were weighed and measured. The 24-hour recall data was analyzed to determine participants' consumption of whole milk, 2% milk, and 1% milk. The milk consumption data was used to calculate grams of milk fat consumed. The cross-sectional association between milk fat intake and severe obesity (BMI  $\geq$  99th percentile) was determined using multivariable logistic regression. Data were available for 145 children, of whom 17% were severely obese. Severely obese children had a lower mean intake of milk fat (5.3 g vs. 8.9 g) and fewer drank any milk (79% versus 95% for not severely obese children ( $p < 0.01$ )). Among the potential confounders assessed, maternal BMI and maternal marital status were associated with severe obesity and were included in a multivariate model. In the multivariate model, higher milk fat consumption was associated with lower odds of severe obesity (OR 0.88 CI 0.80–0.97). Higher milk fat consumption is associated with lower odds of severe obesity among Latino preschoolers. These results call into question recommendations that promote consumption of lower fat milk.

© 2017 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

### 1.1. High fat dairy consumption and obesity

The American Academy of Pediatrics (AAP) and the American Heart Association (AHA) recommend that children ages 2 and older consume skim or 1% milk rather than 2% or whole milk (Gidding et al., 2005, 2006; Daniels et al., 2008). These recommendations are justified by the fact that skim and 1% milk contain fewer calories and less saturated fat than 2% and whole milk, a nutrient profile that is thought to be protective against obesity and cardiovascular risk. The recommendation that children consume 1% or skim milk has broad impacts on nutrition programs serving low-income children in the United States such as the Special Supplemental Nutrition Program for Woman, Infants and Children (2014) (WIC) and school and child care meal programs (2016), all of which promote consumption of low-fat or skim milk.

However, evidence supporting the benefits of lower fat milk over full fat milk is lacking.

Recent research has questioned the link between saturated fat consumption and long-term cardiovascular risk in adults (Chowdhury et al., 2014; Siri-Tarino et al., 2015). Furthermore, recent studies in adults have found that high fat dairy consumption is actually associated with a lower risk of both obesity and diabetes mellitus (Kratz et al., 2013; Holmberg and Thelin, 2013; Rautiainen et al., 2016). However, few studies have been published that examine the association of higher vs. lower fat milk on children's weight and metabolic health. The small number of studies examining this question have either found no association between full-fat milk consumption and weight (O'Connor et al., 2006; Berkey et al., 2005) or have found that full-fat milk consumption is protective against excess weight (Huh et al., 2010; Scharf et al., 2013; Bigornia et al., 2014; Vanderhout et al., 2016). Huh et al. found that higher intake of whole milk at age 2 was associated with a slightly lower BMI-z score at age 3 with no such association found for reduced fat milk (Huh et al., 2010). Similarly, a study of participants in the Early Childhood Longitudinal Study found that children who consumed 2% or whole milk were less likely to be overweight at ages 2 and 4 than those who consumed skim or low-fat milk (Scharf et al., 2013). In addition, Bigornia et al. found that 10 year olds in the highest quartile of full-

\* Funding Source: This study was funded by NIH DK080825, the Hellman Family Foundation, and the NASPGHAN Foundation, NICHD 1K23HD080876-01A1.

☆☆ Conflict of Interest: The authors have no conflicts of interest.

\* Corresponding author.

E-mail addresses: BeckA@peds.ucsf.edu (A.L. Beck), mel.heyman@ucsf.edu (M. Heyman), Cewin.Chao@ucsf.edu (C. Chao), janet.wojcicki@ucsf.edu (J. Wojcicki).

fat milk intake had lower total body fat mass at age 13 compared to those in the lowest quartile (Bigornia et al., 2014). Finally, a recent cross-sectional study of Canadian children ages 12 to 72 months found that consumption of whole milk (vs. 1% or skim) was associated with lower BMI-z score (Vanderhout et al., 2016).

### 1.2. Latinos and obesity

Latino children in the United States have higher rates of obesity than non-Hispanic white children (Ogden et al., 2014), and this disparity emerges before 5 years of age (Rendall et al., 2012). Latino children are also particularly susceptible to severe obesity (Lo et al., 2014a, 2014b) which carries an even greater risk of metabolic sequelae (Lo et al., 2014b; Li et al., 2016; Freedman et al., 2007). Low-income children are also at greater risk of obesity (Singh et al., 2008; Singh et al., 2010) compared to children from higher income families. Thus, understanding how nutritional recommendations and policies may be impacting the risk of severe obesity in low-income Latino children is critical. However, few published studies have evaluated the link between milk fat consumption and weight in low-income Latino children. Consequently, we sought to determine the association of milk fat consumption with weight in low-income, Latino three-year-olds living in San Francisco, California.

## 2. Methods

### 2.1. Study design

We conducted a cross-sectional analysis of data from a birth cohort study of Latino children and mothers to evaluate the association between milk fat consumption and weight at child age 3 years. Please see Wojcicki et al. (2011) for full details of this cohort study.

### 2.2. Participant recruitment and eligibility

Latina mothers were recruited during prenatal visits at two San Francisco Hospitals: San Francisco General and the University of California San Francisco. Mothers were eligible if they self-identified as Latina and were English or Spanish-speaking. Mothers were excluded if they had any of the following medical conditions: drug or alcohol abuse, pre-gestational diabetes or gestational diabetes requiring insulin treatment, polycystic ovarian disease, anorexia nervosa or bulimia, or if they had a health problem that they anticipated would prevent them from breastfeeding. Infants were excluded if they had special health care needs or an Apgar score of <7 at 5 min.

### 2.3. Procedures

At child age 3 years, mothers and children met in-person with a bilingual research assistant from the study. Children were weighed and measured as previously described (Wojcicki et al., 2011). Mothers were also weighed and measured. Child dietary intake was determined via a single 24-hour dietary recall using a multiple-pass method performed by a trained research assistant. All data were collected between May 2009 and June 2010. Dietary intake data were entered into the Food Processor dietary intake analysis program (ESHA Research, Salem, Oregon 2012, version 10.12.0) by an experienced dietary technician. The Food Processor program was used to determine total caloric intake and total fat intake in 24 h for each participant. The dietary technician reviewed each recall to determine the child's total daily intake in ounces of whole milk, 2% milk, 1% milk and skim milk. After the milk data for each dietary recall was entered, there was quality assurance conducted on each recall to confirm that the entered data matched the raw data. The value for ounces of each type of milk consumed was multiplied by the fat content in grams/ounce of the type of milk (whole, 2%, 1% and skim milk) to determine milk fat in grams. Flavored milk was not

included in the analysis due to added sugar content which is likely to have independent effects on adiposity.

### 2.4. Assessment of covariates

At study entry the following variables were assessed via an in-person interview by a bilingual research assistant: highest educational level attained, marital status, country of origin, years in the United States, and preferred language.

### 2.5. Statistical analysis

Maternal and child BMI were calculated and child BMI percentile was determined using the Center for Disease Control (CDC) growth charts. The association between milk fat consumption in grams and child weight status was analyzed using logistic regression models. Milk fat consumption in children was not normally distributed. We first conducted bivariate regression analyses to determine whether there was an association between total grams of milk fat consumed and the following weight categories: 1) BMI  $\geq$  85th percentile (overweight, obese, and severely obese), BMI  $\geq$  95th percentile (obese and severely obese), and BMI  $\geq$  99th percentile (severely obese) (Freedman et al., 2007). We only found a significant association between milk fat consumption and severe obesity. Thus, our subsequent analyses focused on comparing severely obese children (BMI  $\geq$  99th percentile) to those who were not severely obese (all children with BMI < 99th percentile). To determine which covariates to include in our models, we first conducted bivariate logistic regression looking at the association of each of the following potential confounding demographic variables with the outcome of severe child obesity: child gender, maternal BMI, maternal education level (less than or greater than a high school education), maternal marital status, mother's preferred language, and mother's total years in the United States. We also examined the following potential nutritional confounders: total calories, total fat, and total ounces of milk consumed. Significant covariates ( $p < 0.05$ ) were retained in the final multivariate model. *Stata Statistical Software: Release 12*. College Station, TX: StataCorp LP was used for all analyses.

We used descriptive statistics to determine the percentage of all children, severely obese children, and non-severely obese children who had consumed any milk as well as any whole, 2%, 1% or skim milk. We used a chi-squared test to determine the significance of differences in milk consumption patterns between severely obese and not severely obese children. We used simple means to compare milk-fat intake between severely obese and non-severely obese children and used a *t*-test to determine the significance of this difference.

## 3. Results

There were 201 mother-infant dyads enrolled in this cohort. We had BMI data and a complete 24-h dietary recall for 145 children at the 3-year-old visit or 72.1%. Table 1 presents the characteristics of the participants. In our cohort of children, 56% were normal weight (BMI 5th to <85th percentile), 14% were overweight (BMI  $\geq$  85th percentile to <95th percentile), 13% were obese (BMI  $\geq$  95th percentile to <99th percentile) and 17% were severely obese (BMI  $\geq$  99th percentile). Of the mothers, 93% were born outside the United States and spoke Spanish as a primary language, and 80% had a high school diploma or less. Mean maternal BMI was 28.8. The majority of mothers were single but living with a partner (59%), while only 28% were married.

Table 2 presents data on child milk intake. Nearly all of the children in the study (92%) had consumed some form of milk during the 24 h period. However, children who were severely obese were less likely to have consumed any milk (79% vs 95%  $p = 0.007$ ). With regards to the type of milk consumed, 50% of children consumed 2% milk and 28% of children consumed whole milk. Only 15% of children consumed 1% or skim milk. Severely obese children had a tendency toward consuming

Download English Version:

<https://daneshyari.com/en/article/5723600>

Download Persian Version:

<https://daneshyari.com/article/5723600>

[Daneshyari.com](https://daneshyari.com)