

Original Article

e-Consultation Improves Efficacy in Thoracic Surgery Outpatient Clinics[☆]



Nuria M. Novoa,^{a,*} Maria Teresa Gómez,^a María Rodríguez,^a Marcelo F. Jiménez López,^a Jose L. Aranda,^a Elena Bollo de Miguel,^b Florentino Diez,^b Jesús Hernández Hernández,^c Gonzalo Varela^a

^a Servicio de Cirugía Torácica, Hospital Universitario de Salamanca, Salamanca, Spain

^b Servicio de Neumología, Complejo Asistencial Universitario de León, León, Spain

^c Servicio de Neumología, Hospital Nuestra Señora de Sonsoles, Ávila, Spain

ARTICLE INFO

Article history:

Received 6 February 2016

Accepted 3 April 2016

Available online 28 September 2016

Keywords:

Telemedicine

Videoconferencing

e-Consultation

Interdisciplinary health team

Outpatient clinics

Thoracic surgery

ABSTRACT

Objective: The aim of this study is analysing the impact of the systematic versus occasional videoconferencing discussion of patients with two respiratory referral units along 6 years of time over the efficiency of the in-person outpatient clinics of a thoracic surgery service.

Method: Retrospective and comparative study of the evaluated patients through videoconferencing and in-person first visits during two equivalent periods of time: Group A (occasional discussion of cases) between 2008 and 2010 and Group B (weekly regular discussion) 2011 and 2013. Data were obtained from two prospective and electronic data bases. The number of cases discussed using e-consultation, in-person outpatient clinics evaluation and finally operated on under general anesthesia in each period of time are presented. For efficiency criteria, the index: number of operated on cases/number of first visit outpatient clinic patients is created. Non-parametric Wilcoxon test is used for comparison.

Results: The mean number of patients evaluated at the outpatient clinics/year on group A was 563 versus 464 on group B. The median number of cases discussed using videoconferencing/year was 42 for group A versus 136 for group B. The mean number of operated cases/first visit at the outpatient clinics was 0.7 versus 0.87 in group B ($P=.04$).

Conclusions: The systematic regular discussion of cases using videoconferencing has a positive impact on the efficacy of the outpatient clinics of a Thoracic Surgery Service measured in terms of operated cases/first outpatient clinics visit.

© 2016 SEPAR. Published by Elsevier España, S.L.U. All rights reserved.

La discusión de casos por videoconferencia mejora la eficiencia de la consulta externa de cirugía torácica

RESUMEN

Objetivo: El objetivo de este estudio es analizar si la discusión sistemática de casos frente a la ocasional de casos dudosos, mediante videoconferencia, se relaciona con una mejora en la eficiencia de la consulta externa presencial de un servicio de cirugía torácica.

Método: Estudio retrospectivo y comparativo de los pacientes valorados por videoconferencia y consulta externa presencial en 2 periodos de tiempo equivalentes: grupo A (discusión ocasional de casos) entre 2008 y 2010, y grupo B (discusión sistemática semanal) entre 2011 y 2013. Los datos se han extraído de 2 bases de datos prospectivas e informatizadas. Se analizaron el número de pacientes evaluados por videoconferencia y en consulta presencial como primera consulta, y el número de pacientes intervenidos no urgentes bajo anestesia general en cada año y por cada periodo. Se creó el índice casos operados/casos nuevos vistos en consulta externa para cada año y cada periodo. Los datos se compararon usando el test no paramétrico de Wilcoxon.

Palabras clave:

Telemedicina

Videoconferencia

e-Medicina

Reunión multidisciplinaria

Consultas externas

Cirugía torácica

[☆] Please cite this article as: Novoa NM, Gómez MT, Rodríguez M, Jiménez López MF, Aranda JL, Bollo de Miguel E, et al. La discusión de casos por videoconferencia mejora la eficiencia de la consulta externa de cirugía torácica. Arch Bronconeumol. 2016;52:549–552.

* Corresponding author.

E-mail address: nuria.novoa@usal.es (N.M. Novoa).

Resultados: En el grupo A se vieron una media de 563 pacientes en consulta externa frente a 464 en el segundo periodo. La mediana de casos discutidos por videoconferencia/año fue de 42 en el primer periodo frente a 136 en el segundo. La media del índice de casos operados/casos vistos en consulta externa pasó de 0,7 a 0,87 ($p=0,04$) del primer al segundo periodo.

Conclusiones: La discusión sistemática de casos mediante videoconferencia sí tiene una repercusión positiva sobre la eficiencia de la consulta externa medida en términos de casos operados/casos nuevos valorados al mejorar dicho índice.

© 2016 SEPAR. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

The use of e-consultation as an alternative to in-person medical consultations, or as a means of assessing the referral of patients from primary care to specialized care has been studied from various points of view. At first glance, e-consultation, which is more economical than in-person interviews, appears to reduce outpatient clinic costs,¹ although not all authors have found this to be true²; moreover, e-consultations may facilitate and simplify faster access to a specialist,³ and allow medical care to be provided in areas where the population is widely dispersed,⁴ without compromising quality of care.⁵ However, the impact of e-consultation on efficacy in specialist outpatient clinics has not been studied to date.

In May 2006, we began to occasionally use clinician–clinician e-consultation to discuss NSCLC cases with different hospitals within our referral area. Our sporadic experience over a 5-year period was positive, so in 2011 we initiated a program of weekly clinician–clinician e-consultations with 2 hospitals within the referral area. In these multidisciplinary e-consultations,⁶ members of the departments of respiratory medicine, medical oncology and radiotherapy, radiology, and pathology presented the clinical features and radiological images of certain patients to the group and discussed the most suitable treatment.

The aim of this study was to evaluate if systematic clinician–clinician e-consultation versus occasional e-consultation for complicated cases improved efficiency (see definition below in the methods section) in the outpatient clinics of the thoracic surgery department.

Materials and methods

The study was designed as a retrospective, comparative analysis of newly-evaluated NSCLC patients in the outpatient clinics of the department of thoracic surgery over the 6-year study period. Between 2008 and 2010, the department held e-consultations with any departments that requested our opinion on complicated cases. In 2011, we systematized procedures by setting up weekly e-consultations with the respiratory medicine departments of the Hospital Nuestra Señora de Sonsoles de Ávila and the Healthcare Complex of the University of León. These units were selected because they had reliable access to the necessary technical equipment.

Two study groups were established, according to the frequency of the discussions, each comprising a 3-year period: Group A, 2008–2010, years during which occasional discussions were held, and Group B, 2011–2013, years during which regular weekly discussions were held.

Study population

All patients evaluated by e-consultation and first-time in-person interviews in the outpatient clinic during the study period were included. The number of patients operated per year and per study period were also computed; operated cases were defined as all those who underwent scheduled surgery with general anesthesia.

Data were obtained from 2 prospective databases maintained in the department computer system. The first of these lists and classifies all healthcare activity in the unit from the first visit to all subsequent procedures performed in the patient. The second database was created specifically to monitor clinician–clinician e-consultations. This database recorded all cases evaluated by e-consultation, including the date of the discussion, the referring hospital, the number of cases evaluated, the reasons why the decision for the patient was pending, and any reasons for ruling out surgery.

E-consultation equipment

The equipment required for e-consultations included a computer with microphone and webcam together with a corporate application provided by the regional healthcare system (Sacyl) that gave all clinicians participating in the e-consultations access to each others' computer desktop.

Outpatient consultation efficiency parameters. Statistical analysis

The number of patients evaluated by e-consultation and in person, and the number of patients operated by year and by period was calculated.

The ratio of operated cases to new cases seen in the outpatient clinic per year over each time period (2008–2010 and 2011–2013) was calculated. The mean for each 3-year period was compared using Wilcoxon non-parametric statistics. Data analysis was performed using the Stata SE 12.1 statistical package (StataCorp, Texas, USA).

Results

Between 2008 and 2013, a total of 3027 patients were evaluated in a preliminary in-person visit to the outpatient clinic of our department. During this same period, 553 cases were discussed using the clinician–clinician e-consultation system.

The mean number of cases seen in a first visit between 2008 and 2010 was 563/year (range: 518–565). During this period, the ratio of operated cases to new cases/year seen in the outpatient clinic was 0.69, 0.71, and 0.71, respectively (mean: 0.70). The mean number of cases discussed by e-consultation was 42/year (range: 36–60) during these 3 years.

In the period 2011–2013, the mean number of new cases seen in the outpatient clinic was 464/year (range: 442–475). The ratio of operated cases to new cases/year seen in the outpatient clinic during this period was 0.84, 0.89, and 0.87 respectively (mean: 0.87). The mean number of cases evaluated by e-consultation was 136/year (range: 132–147) during these 3 years.

Differences in the mean ratio of operated cases to new cases seen in the outpatient clinic in each period were statistically significant (Wilcoxon $P=0.04$). Fig. 1 shows the number of cases evaluated by e-consultation and the ratio of operated cases to new cases/year seen in the outpatient clinic multiplied by 100 for the purposes of obtaining data that can be reproduced on the same graph.

Download English Version:

<https://daneshyari.com/en/article/5724375>

Download Persian Version:

<https://daneshyari.com/article/5724375>

[Daneshyari.com](https://daneshyari.com)