



CASE REPORT

An unusual cause of acute abdomen— Gas-forming liver abscess due to *Salmonella enteritidis*



Harry Hok Tee Yu, Simon Tsang, Tan To Cheung*

Department of Surgery, University of Hong Kong, Hong Kong, China

Received 23 January 2013; received in revised form 25 March 2013; accepted 9 July 2013
Available online 23 August 2013

KEYWORDS

acute abdomen;
free gas;
gas forming;
liver abscess;
Salmonella enteritidis

Summary Gas-forming pyogenic liver abscess (GFPLA) is considered to be a very severe form of PLA and carries a high mortality. *Klebsiella pneumoniae* is the most common pathogen responsible for the disease, whereas cases where *Salmonella* is cited as the cause are very uncommon. We report the first case of a 53-year-old lady suffering from GFPLA due to *Salmonella*, who was successfully treated with surgical drainage. To the best of our knowledge, this is the first case of GFPLA caused by *Salmonella enteritidis* to be reported in the English literature. Copyright © 2013, Asian Surgical Association. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Pyogenic liver abscess (PLA) is the most common form of liver abscess. Its incidence varies from 1.1 per 100,000 individuals in Europe to 17.6 per 100,000 in Asia. A rising prevalence rate in liver abscess is also reported.¹ The incidence of gas-forming PLAs (GFPLAs) accounts for 10–20% of all PLAs.

Common presentations of PLA include abdominal pain, fever, chills, nausea and vomiting, and a general feeling of

illness. However, the classical triad of abdominal pain, fever, and chills may only account for a third of patients.¹ Overall, 32.5% of GFPLA patients may present with shock, compared to 11.7% in the nongas-forming group.²

Because the mortality of patients suffering from GFPLA can be as high as 27.7% as compared with the 14.4% mortality in the nongas-forming group,² prompt recognition of the condition upon presentation is crucial. Any delay in treatment may result in hospital mortality.

2. Case report

A 53-year-old woman, who works as a waitress in a restaurant, was admitted through the emergency

* Corresponding author. Department of Surgery, University of Hong Kong, Queen Mary Hospital, 102 Pokfulam Road, Hong Kong, China.

E-mail address: tantocheung@hotmail.com (T.T. Cheung).

department for fever and epigastric pain radiating to the back. Apart from osteoarthritis of the knees, which was relieved by simple analgesia, she has enjoyed good health in the past.

An examination on admission revealed epigastric tenderness, dyspnea, and decreased breath sound on the right. Her white blood cell count was $23 \times 10^9/L$ on admission, and the serum bilirubin level was $45 \mu\text{mol/L}$. Other laboratory results were as follows: serum alkaline phosphatase level, 132 U/L; serum aspartate aminotransferase level, 328 U/L; serum alanine aminotransferase, 376 U/L; international normalized ratio, 1.8; serum glucose level, 1.7 mmol/L.

Radiography of the chest was performed (Fig. 1). The chest X-ray showed mild haziness over the right lower zone of the lung. In addition, there was a lucency occupying the right subphrenic area. In view of the patient's clinical condition and abnormal chest X-ray result, an urgent computed tomography (CT) scan of the abdomen with contrast was performed that showed a 16.5-cm gas-forming liver abscess involving segments 6–8 (Fig. 2).

CT-guided drainage of the liver abscess was performed, but only a minimal amount of pus was obtained. The patient continued to run a high fever with hypotension. A decision was made to perform laparotomy and open drainage of the liver abscess. During the laparotomy, a large amount of turbid fluid was noted in the peritoneal cavity. The liver was grossly edematous and the large abscess was seen at the dome of the liver with pus and gas discharge. Open drainage was then performed. Operative cholangiogram showed contrast leakage into the abscess cavity, indicating the communication of the bile duct to the abscess. The common bile duct was explored, and a T-tube was placed to facilitate biliary drainage in a controlled manner (Fig. 3). Abdominal drains were placed in the abscess cavity and the subhepatic area. The patient was supported in the intensive care unit after the operation. Both abscess tissue and pus only yielded *Salmonella*




Figure 1 Erect chest X-ray on admission: lucency below the right diaphragm.




Figure 2 Contrast computed tomography: faintly rim-enhancing gas-forming abscess in the right lobe of the liver.

enteritidis. The bacteria were sensitive to piperacillin. A histology section of the abscess did not yield any evidence of malignancy. The abdominal drains were removed 3 weeks after the operation. The patient was given 3 weeks of intravenous piperacillin and 3 weeks of oral amoxicillin/cavulanate, and was discharged home 49 days after the procedure. The T tube was removed 6 weeks after the operation. A reassessment contrast CT scan showed resolution of the liver abscess at 3 months after the operation (Fig. 4).

3. Discussion

GFPLA is a rare condition that carries a high mortality. The usual cause of PLA is mainly due to portal seeding from appendicitis and diverticulitis. With the improvement of antibiotics, biliary tract diseases have become the most common identifiable cause, although most of them are cryptogenic in origin.³ Patients with compromised immunology may be more often affected. Diabetes mellitus is not regarded as a cause of PLA; however, a high prevalence of the disease is noted among patients suffering from PLA in Asia. Reports suggested that about 43% of patients suffering from PLA are known to have diabetes.^{1,4} An even higher proportion of diabetic patients are noted in gas-forming entities associated with higher mortality rates.^{1,2,4}

In the management of this patient, a prompt decision to operate was made. Apart from drainage of the liver abscess, cholecystectomy was performed as cholecystitis was the usual source for liver abscess formation. Operative cholangiogram was performed in the same operation to identify the presence of a possible biliary stone or sludge that could have led to liver abscess formation. Because the liver abscess contained a large portion of solid debris, open drainage and debridement of the abscess had been performed. The displaced intrahepatic duct was unavoidably broken down together with the abscess. Because the

Download English Version:

<https://daneshyari.com/en/article/5731487>

Download Persian Version:

<https://daneshyari.com/article/5731487>

[Daneshyari.com](https://daneshyari.com)