



ELSEVIER

Contents lists available at ScienceDirect

## International Journal of Surgery Case Reports

journal homepage: [www.casereports.com](http://www.casereports.com)

# Management of the open abdomen using negative pressure wound therapy with instillation in severe abdominal sepsis

## A review of 48 cases in Hospital Mexico, Costa Rica

Pablo Sibaja<sup>a,b,\*</sup>, Alfredo Sanchez<sup>a</sup>, Guillermo Villegas<sup>a,b</sup>, Alvaro Apestegui<sup>b</sup>, Esteban Mora<sup>b</sup>

<sup>a</sup> Universidad San Judas Tadeo, San Jose, Costa Rica

<sup>b</sup> Hospital Mexico, La Uruca, Costa Rica

## ARTICLE INFO

## Article history:

Received 11 October 2016

Accepted 15 November 2016

Available online 17 November 2016

## Keywords:

Intra-abdominal sepsis

Negative pressure wound therapy

Negative pressure wound therapy with instillation

Open abdomen

## ABSTRACT

**INTRODUCTION:** Despite the numerous advances in recent years, severe abdominal sepsis (with associated organ failure associated with infection) remains a serious, life-threatening condition with a high mortality rate. OA is a viable alternative to the previously used scheduled repeat laparotomy or continuous peritoneal lavage. The use of Negative Pressure Wound Therapy (NPWT) has been described as a successful method of management of the open abdomen. Adding instillation of saline solution to NPWT in a programmed and controlled manner, could offer the clinician an additional tool for the management of complex septic abdomen.

**OBJECTIVES:** To explore if the concept of active two-way therapy (Negative pressure wound therapy with instillation or NPWT-I) yields superior control of underlying, life-threatening abdominal infections and its effects on survival and morbidity in patients with severe abdominal sepsis when management with an open abdomen is required.

**METHODS:** A retrospective review of 48 patients with severe abdominal sepsis, who were managed with and open abdomen and NPWT-I was performed. NPWT-I was initiated utilizing the same parameters on all patients, this consisted of cycles of instillation of saline solution, which was removed through negative pressure after a short dwell period. We observed the effects on primary fascia closure rate, mortality, hospital and SICU length of stay and associated complications.

**RESULTS:** Our patient group consisted of 20 (42%) males and 28 (58%) females. Average age was 48 years. Mortality in these patients was attributed to pulmonary embolism (n = 1), acute renal failure (n = 2) and cardiopulmonary arrest (n = 1). Average total hospital stay was 24 days, and stay in the SICU (n = 26) averaged 7.5 days.

No acute complications related to the NPWT-I. All patients presenting with abdominal compartment syndrome resolved after initiation of the NPWT-I. A total of 46 patients (96%) patients achieved fascia closure after NPWT-I therapy after an average of 6 days. Four patients (8%) died during the course of treatment of causes unrelated to NPWT-I.

**CONCLUSION:** This therapy showed added benefits when compared to traditional methods such as "Bogota bag, Wittmann patch, or NPWT traditional in the management of the open abdomen pertaining to severe abdominal sepsis.

NPWT-I in patients with severe abdominal sepsis had promising results, since we obtained higher fascia closure rates, lower mortality and reduced hospital and ICU length of stay with no complications due to this therapeutic approach.

© 2016 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### 1. Introduction

Despite numerous advances in recent years, severe abdominal sepsis (organ failure associated with infection) remains a serious, life-threatening condition with a high mortality rate. Because of

breakthroughs in trauma care that require damage control surgery, the use open abdomen (OA) management has increased in scope and complexity.

In non-coronary ICU's, severe sepsis (SS) is considered the main cause of death. In the USA alone, 900,000 cases of SS are diagnosed every year and it causes 210,000 deaths every year. In Europe, the number of deaths per year is 150,000 [1,2].

Peritonitis remains the main cause of developing SS, and up to 11% of patients admitted to the surgical ICU progress to this state

\* Corresponding author at: Universidad San Judas Tadeo, San Jose, Costa Rica.  
E-mail address: [drsibaja@usanjudas.ac.cr](mailto:drsibaja@usanjudas.ac.cr) (P. Sibaja).

due to a previous intra-abdominal infection [3]. Currently, mortality rates for intra-abdominal sepsis ranges between 30% and 40% [4,5].

During the past two decades, there has been a paradigm shift toward management of patients with severe abdominal sepsis by OA as a viable alternative to the previously used scheduled repeat laparotomy or continuous peritoneal lavage [6], as to deal with or prevent recurrent infection. In addition, it is well established that the visceral or retroperitoneal edema secondary to shock and reperfusion, may increase intra-abdominal pressure to dangerous levels, leading intra-abdominal hypertension and organ dysfunction [7]. Patients with this constellation of symptoms must have their abdomens left temporarily open to allow for visceral and renal perfusion as well as adequate pulmonary function.

Negative pressure wound therapy (NPWT) is one of a variety of techniques used to manage the OA. NPWT was initially introduced in 1996 [8], with the intention of treating chronic soft tissue wounds. Later, it found its way into the treatment options of open abdomen. NPWT with Instillation (NPWT-I) has become a variant of this method. Although this technique was not intended for treatment of open abdomen, its potential as a new therapeutic tool, incited us to use it in patient with a septic open abdomen, and we have continued utilizing this technique thereafter.

In this report, we describe our use of negative pressure wound therapy as delivered by V.A.C.<sup>®</sup> Therapy (KCI USA, Inc., San Antonio, TX) in addition to timed, automated instillation of saline in the management of patients with SS.

## 2. Overview

Historically, the OA was treated with simpler approaches such as “Bogota bag” [9], Wittmann Patch [10] and Barker’s vacuum Pack [11], which yielded a variety of complications [12] such as marked adhesion formation, development of enteric fistula, non-quantifiable loss of fluids, evisceration, hemorrhage, contamination of the abdominal cavity surgical wound (especially when in proximity to stomas), spread of bacteria into the ICU and ward environment, and a high rate of subsequent ventral hernias. Different methods of temporary abdominal closure (TAC) have been developed to protect the temporary open abdomen and decrease complications [12,13].

NPWT resulted in greater rates of fascia closure [14,15] obviating the need for subsequent hernia repair in many patients [16]. The utility of this technique is not limited to the early postoperative period, but can be successful in for up to 3–4 weeks after the initial operation [17]. Recent large scale studies have reinforced the benefits of NPWT as compared to other TAC methods [18], and it’s early application has been showed to be beneficial [17].

Peritoneal negative pressure therapy (PNPT) decreases systemic inflammation and organ damage [19]. PNPT served to evacuate a significantly greater volume of ascites than passive drainage. Systemic inflammation as measured by serum levels of TNF-I, IL-1 and IL-6, was significantly reduced in the PNPT group and was associated with significant improvement in histology of the intestine, lung, kidney, and liver [20]. Efficacy was attributed in part to an attenuation of peritoneal inflammation by the removal of cytokines and other biochemical mediators in the ascitic fluid [19]. Interruption of peritoneal hemolymphodynamics via controlled vacuum aspiration appears to decrease entry of inflammatory mediators into the blood stream, thereby possibly limiting systemic inflammation [21]. The clinical implication of these studies was that because sepsis/trauma results in an inflammatory ascites that may perpetuate organ injury; removal of the ascites can break the cycle and decrease organ damage via source control [19].

The primary goal of TAC is to create a tension-free closure of the abdomen without increasing intra-abdominal pressure. Attempting primary fascia closure under tension in patients with extensive abdominal wall and intra-abdominal organ edema, is associated with an increased incidence of multiple organ failure (MOF), necrotizing abdominal wall infections, and mortality [7]. The optimal method of TAC should contain and protect the contents of the peritoneal cavity from external contamination and injury, preserve fascia; minimize desiccation and damage to viscera, remove and quantify third space fluid; prevent loss of domain, lower bacterial count, inflammatory response, keep patient’s abdominal wall skin dry and intact; preserve the integrity of the abdominal wall, be simple to perform and maintain, provide ease of reentry and have minimal adverse physiologic effects [13,22,23]. Various TAC methods have been subject to multiple studies, and their advantages and disadvantages are known [24], but are too numerous to cite. Although no prospective randomized studies are available to compare effectiveness of various TAC techniques as compared to NPWT-I, some evidence exists that shows a beneficial effect of this technique in the management of complex abdominal pathology [25].

## 3. Methods

A retrospective review was performed of 48 patients with clear indication for open abdomen management due to severe abdominal sepsis that were treated with NPWT-I between November, 2007 and November, 2008 by the SICU team of the Hospital Mexico, in San Jose, Costa Rica. The following criteria were utilized in the selection of patients who would receive this treatment: Björck open abdomen class 2b or greater [24], patients with an APACHE score of 12 or greater at admission with a diagnosis of abdominal sepsis, with contamination or with secondary generalized peritonitis. All other patients that required management with an open abdomen who did not meet the previously expressed criteria were excluded from our study.

The scope of our study involves only the use of NPWT-I in regards to the management of intra-abdominal sepsis, other considerations such as abdominal closure, management of abdominal compartment syndrome and other associated conditions are not fully address and are not the main consideration of this paper.

Our study group consisted of 20 males and 28 females, with ages between 22 and 76 years, in whom the infectious source was found but not controlled during the first intervention and management via OA and abdominal cavity washout for residual contamination was scheduled. All patients selected received NPWT-I. The procedure was performed by applying a sterile abdominal dressing, which consists of a fenestrated soft plastic non-adherent layer with enclosed central foam, which is placed on the surface of the viscera. Then, two layers of V.A.C. GranuFoam<sup>®</sup> dressings are applied over the plastic layer, the first layer in direct contact with the non-adherent plastic and located inside the abdominal cavity under the edges of the incision and the second, is placed directly in the abdominal incision filing the exact space of the wound opening (Fig. 1). Finally, a transparent adhesive is placed over the GranuFoam<sup>®</sup> and the wound to seal the abdominal cavity. A 1-inch diameter orifice is cut off the plastic seal on the lower or caudal area of the wound to connect a Sensa T.R.A.C.<sup>®</sup> adhesive pad, which will serve as the outflow tubing, that will be later connected from the dressing to the canister in the negative pressure device. Before the final connection is made, test of the integrity (seal) of the vacuum system must be performed by connecting a sterile suction tube to the wound system, and thus guaranteeing that no leaks are present. Finally, a second 1-inch dressing is placed over the wound to adhere the instillation tubing, which is later connected to a saline

Download English Version:

<https://daneshyari.com/en/article/5732567>

Download Persian Version:

<https://daneshyari.com/article/5732567>

[Daneshyari.com](https://daneshyari.com)