



Contents lists available at ScienceDirect

## International Journal of Surgery Case Reports

journal homepage: [www.casereports.com](http://www.casereports.com)

# Total proximal anastomosis detachment after classical Bentall procedure



Aref Rashed\*, Karoly Gombocz, Andras Vigh, Nasri Alotti

Department of Cardiac Surgery, Zala St. Raphael County Hospital, Zalaegerszeg, Hungary

## ARTICLE INFO

### Article history:

Received 2 February 2017  
Received in revised form 5 June 2017  
Accepted 11 June 2017  
Available online 27 June 2017

### Keywords:

Bentall procedure  
Case report  
Left ventricle outflow tract elongation  
Pseudoaneurysm

## ABSTRACT

**INTRODUCTION:** Since its introduction in 1968, the Bentall procedure has been the primary surgical solution for aneurysms of the aortic root. However, many surgeons have reported serious procedural complications such as detachment of coronary ostia and pseudoaneurysm formation at anastomosis sites. Therefore, the Bentall procedure has undergone several modifications to eliminate those complications. Partial or total detachment of the proximal anastomosis is rarely reported.

**PRESENTATION OF CASE:** We report a total detachment of the proximal anastomosis after a Bentall operation with emphasis on the possible practical mechanisms, which might have led to the development of this very rare complication. The diagnosis was confirmed at a routine follow up examination and urgent surgery was performed. We also report our operative solution and review other possible surgical solutions that might be considered in this setting.

**DISCUSSION:** The Bentall procedure and its modifications continue to be considered the gold standard for treating aneurysms involving the aortic root. Various modifications can serve as optimal solutions for procedure-related complications.

**CONCLUSION:** Surgeons performing the Bentall procedure must be familiar with all existing modifications because they are complementary to the original surgical procedure. In the absence of endocarditis left ventricle outflow tract elongation may be an acceptable surgical solution to deal with total detachment of the proximal anastomosis.

© 2017 The Author(s). Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).


## 1. Introduction

Since its introduction in 1968, the Bentall procedure has been the primary surgical solution for aneurysms of the aortic root [1]. Due to procedural complications such as bleeding, detachment of coronary ostia and pseudoaneurysm formation at anastomosis sites, the Bentall procedure has undergone several modifications [2–4]. Today surgeons typically perform modified Bentall operations instead of the classic Bentall operation, which involves direct coronary anastomosis and wrapping the aneurysmal aortic wall around the valved conduit. Detachment of the proximal suture line after the Bentall procedure is a rarely reported [9,10]. Here, we report a total detachment of the proximal anastomosis after a Bentall operation induced stress on the possible mechanisms, which might have led to development of this very rare complication. We also report our surgical solution in dealing with this frightening situation. This work has been reported in line with the SCARE criteria [5].

## 2. Presentation of the case

A 42-year-old man with Marfan syndrome underwent a Bentall procedure due to a 6 cm aortic root aneurysm and severe bicuspid aortic valve regurgitation. In his past history aortic coarctation and hypertension were highlighted. Due to aortic coarctation he underwent a left thoracotomy with excision of the coarctic segment and end-to-end anastomosis when he was 7 months old. At the age of 17, a bypass with prosthetic graft was anastomosed between the left subclavian artery and the ascending aorta due to recoarctation. During routine follow up, echocardiography revealed a dilated aortic root and significant aortic valve regurgitation. Computer tomography (CT) images showed a root aneurysm 6 cm in diameter and minimal hypoplasia of the aortic arch. Coronarography revealed a normal coronarogram. Surgery was performed via mid-line sternotomy and extracorporeal circulation with cannulation of the proximal aortic arch and right atrium. The Bentall procedure was performed using a home-made mechanical valve-graft conduit (a 27-mm Sorin bileaflet mechanical valve sutured to the end of a 34-mm Polythese ICT vascular prosthesis using continuous 2/0 prolene suture), which was sewn to the aortic ring using supra annular interrupted 2-0 mattress pledgeted sutures. The coronary ostia were reimplemented applying the button technique, and

\* Corresponding author at: Department of Cardiac Surgery, Zala St. Raphael County Hospital, Zrínyi M. str. 1., Zalaegerszeg 8900, Hungary.  
E-mail address: [aref.rashed.szv@zmkorhaz.hu](mailto:aref.rashed.szv@zmkorhaz.hu) (A. Rashed).




**Fig. 1.** Totally detached valved conduit with empty aortic ring; LV: left ventricle, IVS: interventricular septum, LA: left atrium, MV: mechanical valve, VP: vascular prosthesis, PS: pseudoaneurysm.

the aneurysmal wall was wrapped over the vascular prosthesis. The postoperative period was uneventful except for early excessive bleeding (1st h: 250 ml, 2nd h: 180 ml and 1050 ml over 24 h), which was treated successfully using a conservative correction of the coagulation parameters.

As a follow-up measure, transthoracic echocardiography (TTE) was performed every 6 months by his cardiologist; according to the available documents, no abnormalities were detected. At the patient's last follow up, 26 months after the operation, he visited his cardiologist and complained of exertional dyspnea. Transthoracic, transesophageal echocardiography and CT scan revealed a pseudoaneurysm around the graft and the mechanical valve was totally detached from the aortic annulus. The distance between the original aortic annulus and the detached mechanical valved composite was approximately 20 mm (Fig. 1).

In an urgent operation, cardiopulmonary bypass was initiated prior to re-sternotomy with cannulation of the femoral vessels. After re-sternotomy and careful adhaesiolysis, the ascending aorta was clamped directly below the origin of the innominate artery and the wall of the pseudoaneurysmal sac was opened. The heart was arrested by direct administration of cold crystalloid cardioplegia into the valved conduit. The valved conduit was found to be hanging above the aortic ring, fully detached and suspended in place by the main coronary arteries. No tension or traction was observed on the coronary arteries. The aortic annulus was empty (Fig. 2). As no signs of endocarditis were observed, we decided to elongate the left ventricle outflow tract (LVOT) using a vascular prosthesis between the original aortic ring and the sewing ring of the valved conduit. A vascular graft with 30-mm in diameter and 2.5- cm in length was selected. The proximal anastomosis, at the level of the original aortic annulus, was prepared using running 3-0 polypropylene sutures with external reinforcement incorporating a strip of Teflon. After removing all the disrupted pledgeted sutures from the ring of the mechanical valve, the other end of the vascular graft was sewn to the ring of the mechanical valve using running 3-0 polypropy-




**Fig. 2.** Operative seen; AR: empty aortic ring, MV: mechanical valve, VP: vascular prosthesis, PSW: pseudoaneurysmal wall.

lene sutures. The aneurysmal sac was left opened, unwrapped and a small drain was placed at its lowest point.

The postoperative period was uneventful. Echocardiographic studies revealed no significant pressure gradient at the level of the supra annular mechanical valve and no kinking of the vascular graft sewed to the aortic ring (Fig. 3). Eight months after the reoperative procedure, the patient was completely asymptomatic.

**3. Discussion**

The Bentall procedure and its modifications have continued to be considered the gold standard for treating aneurysms involving the aortic root [1]. Various modifications can serve as optimal solutions for procedure-related complications. Complications of the Bentall procedure can be divided into two subgroups: endocarditis or thromboembolic events and procedure-related complications. The former is valve-related complications occur-



**Fig. 3.** LVOT elongation with a new short vascular prosthesis; AR: aortic ring, MV: mechanical valve, nVP: new vascular prosthesis.

Download English Version:

<https://daneshyari.com/en/article/5732699>

Download Persian Version:

<https://daneshyari.com/article/5732699>

[Daneshyari.com](https://daneshyari.com)