



Contents lists available at ScienceDirect

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

Severe fatal protein malnutrition and liver failure in a morbidly obese patient after mini-gastric bypass surgery: Case report



Mohammad Ali Kalantar Motamedi^a, Maryam Barzin^a, Mohammadreza Ebrahimi^b,
Reza Ebrahimi^b, Alireza Khalaj^{b,*}

^a Obesity Research Center, Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran

^b Obesity Treatment Center, Department of Surgery, Shahed University, Tehran, Iran

ARTICLE INFO

Article history:

Received 2 January 2017

Received in revised form 18 February 2017

Accepted 18 February 2017

Available online 21 February 2017

Keywords:

Case report

Morbid obesity

Malnutrition

Liver failure

Bariatric surgery

Mini-gastric bypass

ABSTRACT

INTRODUCTION: Mini-gastric bypass (MGB) is a bariatric surgical technique popular in many centers due to shorter duration, easier technique, and excellent weight loss results. However, it may be associated with postoperative malnutrition. This case describes the clinical course and unfortunate outcome of a morbidly obese patient who underwent MGB and developed malnutrition in the first postoperative year. **PRESENTATION OF CASE:** A 37 year-old female patient with a BMI of 44 kg/m² successfully underwent MGB surgery in June 2015 and was discharged uneventfully. She presented with lower extremity edema and generalized weakness 8 months later, with a blood albumin level of 3.1 g/dL, compared to a normal preoperative value. She was admitted and received a high-protein diet, and her clinical condition improved. Three months after her discharge, she was readmitted with the same complaints, as well as pancytopenia. She was also hypocupremic. After unsuccessful intensive supportive measures, she finally underwent revisional gastrogastrostomy. However, she developed signs and symptoms of profound liver failure postoperatively (albumin 1.8 g/dL; total bilirubin 7.5 mg/dL; prothrombin time 34 s) and pancytopenia persisted. All resuscitative measures were unsuccessful and she expired in July 2016. **DISCUSSION:** Multiple factors can contribute to postoperative malnutrition and liver dysfunction after MGB, including the presence of baseline liver disease, inadequate diet supplementation, leaving a too-short common small intestinal channel, and ethnic variations in small bowel length. These factors should also be considered when deciding to perform corrective surgery.

CONCLUSION: Careful, individualized treatment and follow-up plans may help to prevent such catastrophic consequences.

© 2017 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Given the widespread magnitude of obesity, bariatric surgery is now a common procedure worldwide. It effectively lowers body weight and resolves obesity-related complications, and is superior to other means of weight loss. However, as with any invasive procedure, bariatric surgical techniques are not without risks.

There are different techniques of bariatric surgery, and mini-gastric bypass (MGB)¹ is a newly developed method. This technique uses a long gastric tube with an antecolic loop gastrojejunostomy. A long gastric tube is created using an Endo GIA[®] stapler approximately 1.5 cm to the left of the lesser curvature from the antrum to the angle of His. Then, a loop gastroenterostomy is created with

the small bowel about 200 cm distal to the ligament of Trietz. MGB is shorter in operative time than conventional Roux-en-Y gastric bypass (RYGB) and also sleeve gastrectomy, with comparable or even better weight loss results [1–5]. However, this method is associated with a greater post-op malnutrition rate of approximately 1% requiring reoperation, due to the longer bypassed limb [5–7]. We hereby report a case of severe malnutrition nearly 1 year after MGB, who presented to us and was managed in an academic setting, in line with the SCARE criteria [8].

2. Presentation of case

Our patient was a 37 year-old Middle Eastern morbidly obese female who presented with a body mass index (BMI) of 44 kg/m² (height 166 cm, weight 115.5 kg). She was referred to our obesity clinic to be considered for bariatric surgery, as part of the Tehran Obesity Treatment Study [9]. Her past history was significant for diabetes since 5 years ago, dyslipidemia, hypothyroidism, and a history of nephrolithiasis. She received oral glycemic

* Corresponding author at: Shahed University, P.O. Box: 18155-159, Tehran, Iran.
E-mail address: khalaj@totc.ir (A. Khalaj).

¹ MGB – mini-gastric bypass, RYGB – Roux-en-Y gastric bypass, BMI – body mass index.

Table 1
Pre- and postoperative weight, hematologic, and biochemical indices after mini-gastric bypass and corrective surgery.

Variables	Normal values	Course of progression							
		Mini-gastric bypass					Corrective surgery		
		Preoperative	Postoperative					Postoperative	
		1 mo	3 mo	6 mo	9mo	12 mo	12.5 mo	13 mo	
Weight (kg)		115.5	104.1	84.7	76.2	72.8	63	–	63.7
BMI (kg/m ²)	25–25	44	39.7	32.3	29	27.8	24	–	24.3
EWL (%)	–	–	22.8	61.7	77.7	–	104.1	–	103
Albumin (g/dL)	3.5–5.2	4.1	–	–	2.3	2.4	1.8	2.9	1.7
Ferritin (ng/mL)	11–307	12.7	–	–	–	–	–	109	–
AST (U/L)	Up to 31	16	–	–	30	31	43	83	29
ALT (U/L)	Up to 31	23	–	–	29	30	60	64	33
ALP (U/L)	100–290	64	–	–	56	70	395	312	78
Total bilirubin (mg/dL)	Up to 1.3	0.7	–	–	–	–	1.1	1.28	7.5
PT (s)	13–Nov	12	–	–	–	11.3	10	16.3	34
Copper (µg/dL)	70–140	103	–	–	62	–	44	–	40
Calcium (mg/dL)	50–170	9.1	–	–	–	8.7	8.3	8.6	8.7
FPG (mg/dL)	70–110	137	91	98	87	85	81	92	94
HbA1c (%)	5–7.5	6.8	6.5	6.1	4.9	4.7	4.8	–	4.6
Red blood cells (10 ⁶ /mL)	4.2–5.4	3.97	–	–	–	–	3.77	2.9	–
Hemoglobin (g/dL)	14–17.5	11.9	–	–	13.4	9.1	8	7.4	8.6
Hematocrit (%)	42–52	36.9	–	–	41	27.1	25.5	25	25.7
MCV (fl)		86.4	–	–	–	–	85.4	86.2	–
Platelets (×1000/micL)	150–450	144	–	–	128	78	55	83	68

BMI; body mass index, EWL; excess weight loss, AST; aspartate aminotransferase, ALT; alanine aminotransferase, ALP; alkaline phosphatase, PT; prothrombin time, FPG; fasting plasma glucose, MCV; mean corpuscular volume.

agents (metformin, pioglitazone) as well as insulin, atorvastatin, levothyroxine, and aspirin daily. She was evaluated by a multidisciplinary team of obesity experts.

Her initial assessment demonstrated a normal pulmonary function test and electrocardiogram. Upper GI endoscopy was normal. Abdominal ultrasound study revealed grade 3 fatty liver with an increased liver span of 15.6 cm.

She underwent mini-gastric bypass with the method described above successfully in June 2015, and was discharged in good health after 3 days. She received routine postoperative follow-up visits by a standard protocol at 1, 3, and 6 months, and various measurements were made at each visit including blood indices. These values are presented in Table 1. She received oral multivitamin-mineral capsules daily postoperatively (Pharmaton®, Boehringer Ingelheim Inc., Ingelheim am Rhein, Germany), which contains 2 mg copper (200% recommended daily allowance (RDA)), 10 mg ferrous sulfate (71% RDA), 100 mg folic acid (50% RDA), and 1mcg vitamin B12 (40%), as well as vitamins A, B group, C, D, E, nicotinamide, biotin, and minerals, calcium, magnesium, zinc, and selenium.

She had lost weight significantly throughout her follow-up (Table 1) and her diabetes was completely resolved. It was not until the 8th month after surgery that she presented with lower extremity edema, weakness, and generalized fatigue. She was admitted to hospital and found to have an albumin value of 3.1 g/dL, as compared to a preoperative value of 4.1 g/dL. She received a high-protein diet and intensive measures and was discharged after two weeks. She was well until 3 months later, at which time she presented again with persistent lower extremity edema despite supportive measures. Her complete blood count showed a platelet count of 55,000/µL, as well as a hemoglobin level of 8 g/dL. Her liver function tests showed slightly increased aspartate aminotransferase (AST) level of 43 U/L, alanine aminotransferase (ALT) level of 60 U/L. She was also found to be hypocupremic (44 µg/dL).

During workup for pancytopenia, a bone marrow biopsy was done, which showed adequate maturation in all cell lines, and was not in favor of any specific diagnosis. Liver biopsy was done and showed non-specific inflammation in portal spaces and fatty change. Corrective surgery was initially postponed to address the thrombocytopenia; however, such surgery was deemed necessary

as the patient was deteriorating and not responding to supportive measures such as erythropoietin administration. Her prothrombin time (PT) rose to 16.3 s and total bilirubin to 1.28 mg/dL (direct bilirubin 0.71 mg/dL).

She underwent gastrogastrostomy 13 months after her first bariatric surgery and nearly four months after showing early signs and symptoms of liver failure. After the surgery, her liver function did not improve, and her albumin level dropped to 1.7 g/dL. All supportive measures failed to reverse this downhill course, and overt liver dysfunction developed with a total bilirubin of 7.5 mg/dL and a PT of 34 s, extensive ecchymosis, dyspnea, and multi-organ failure. She expired in July 2016. Her autopsy examination revealed a common intestinal channel of 108 cm.

3. Discussion

Although RYGB is considered the gold standard of bariatric surgery for many years, MGB, a relatively new modification of loop gastric bypass first described by Rutledge in 2001 [10], is popular in some centers due to multiple benefits such as shorter operative times, easier technique, and impressive weight loss results [6,10]. However, besides from the inherent risk of bile reflux, marginal ulcers, and long-term risk of gastric cancer, malnutrition is a major concern. Kruschitz et al. recently published their results [11] of comparison between MGB and RYGB regarding liver parameters after surgery in 50 patients. They found significantly poorer liver function in patients undergoing MGB in the first year not associated with weight loss, a finding that may necessitate a more intensive postoperative care in MGB patients. However, in our experience, this procedure has been performed successfully in our center for a couple of years with excellent weight loss results and correction of obesity-related complications, and this is our first and only unfortunate outcome.

The authors believe this case report highlights the importance of a more individualized approach to bariatric surgery and MGB. Selection of a 200 cm efferent limb may be suitable for most, but not for all patients. An alternate and safer approach may be careful inspection of each case to assure not to leave a too-short common channel to the ileocecal valve, as there may be differences in the

Download English Version:

<https://daneshyari.com/en/article/5732869>

Download Persian Version:

<https://daneshyari.com/article/5732869>

[Daneshyari.com](https://daneshyari.com)