

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

Acute abdomen in a patient with paraesophageal hernia, resulting in acute compromised respiratory function: A case report



Mira Mekhael, Alaa El-Hussuna*

Department of Gastrointestinal Surgery, Aalborg University Hospital, 9000 Aalborg, Denmark

ARTICLE INFO

Article history:

Received 5 October 2016

Received in revised form 18 January 2017

Accepted 19 January 2017

Available online 23 January 2017

Keywords:

Paraesophageal hernia

Acute abdomen

Compromised respiratory function

Elective hernia repair

Watchful waiting

Case report

ABSTRACT

INTRODUCTION: We present a case of acute abdomen, causing increased intra-abdominal pressure, leading to further herniation of an existing paraesophageal hernia, and consequently acute compromised respiratory function. This acute respiratory complication to a paraesophageal hernia has not previously been reported.

PRESENTATION OF CASE: We present a case of a 75-year-old female who was acutely admitted with stridor. The patient was known to have a paraesophageal hernia monitored using watchful waiting, and dyspnoea. The patient's condition deteriorated, leading to intubation. Diagnostic imaging revealed a paraesophageal hernia pressing onto the trachea as well as appendicitis and ileus. Surgery confirmed perforated appendicitis, peritonitis, and ileus causing high intra-abdominal pressure, resulting in further herniation of the paraesophageal hernia as a cause for acute compromised respiratory function. Appendectomy and gastropexy were performed. The patient was later discharged to rehabilitation.

DISCUSSION: Patients with pulmonary symptoms caused by a paraesophageal hernia, especially patients with sizeable hernias, could potentially be in greater risk of severe airway affection if complicated by acute abdomen. These patients could benefit from elective hernia repair, rather than watchful waiting, as it would eliminate pulmonary symptoms and prevent similar cases. Patients monitored using watchful waiting should be informed that acute abdomen could cause acute compromised respiratory function.

CONCLUSION: Any case of acute abdomen causing high intra-abdominal pressure could potentially cause further herniation of an existing paraesophageal hernia, resulting in acute compromised respiratory function. In patients known to have a paraesophageal hernia, similar cases should be suspected if the patient presents with acute breathing difficulties.

© 2017 The Authors. Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

A paraesophageal hernia occurs when part of the gastric fundus herniates through the esophageal hiatus and lies alongside the esophagus [1]. Paraesophageal hernias comprise 5% of all hiatal hernias and are more common in the elderly [1,2]. The cause of hiatal hernias is related to increased intra-abdominal pressure, causing a transdiaphragmatic pressure gradient at the gastroesophageal junction. This results in weakening of the phrenoesophageal membrane and widening of the esophageal hiatus [3]. Most patients with a paraesophageal hernia are believed to be asymptomatic. Symptoms can be caused by obstruction, gastroesophageal reflux disease, bleeding, and iron deficiency anaemia. Other non-specific symptoms associated with paraesophageal hernias are postprandial fullness, chest pain, and breathing difficulties [2,3]. Patients

with paraesophageal hernias can also present acutely with gastric volvulus and incarceration, which requires immediate surgical intervention [2,3]. Currently, asymptomatic or minimally symptomatic paraesophageal hernias are monitored using watchful waiting, and only symptomatic hernias are recommended for elective hernia repair [2].

To our knowledge, we report the first case of acute abdomen causing high intra-abdominal pressure, resulting in further herniation of an existing paraesophageal hernia, causing the hernia to press onto the trachea, consequently leading to acute compromised respiratory function.

It is well known that perforated appendicitis can cause peritonitis and ileus [4]. This, and other acute abdominal syndromes such as intestinal perforation, acute pancreatitis or trauma, frequently led to high intra-abdominal pressure [5]. In this case, peritonitis and ileus were the reason for the increased intra-abdominal pressure, but any case of acute abdomen leading to high intra-abdominal pressure could potentially cause cases like this.

It is also known that paraesophageal hernias can cause pulmonary symptoms [2]. However, this case shows that a case of acute

* Corresponding author at: Skovbyvej 18, 2740 Skovlunde, Denmark.

E-mail addresses: mekhael.m@hotmail.com (M. Mekhael), alaa@itu.dk (A. El-Hussuna).



Fig. 1. Thoracoabdominal CT scan showing ileus and appendicitis.

abdomen possibly can cause acute exacerbation of pulmonary symptoms associated with a paraesophageal hernia, resulting in acute compromised respiratory function.

2. Case report

A 75-year-old Caucasian female with a history of depression, atrial fibrillation, dilated cardiomyopathy, a paraesophageal hernia currently monitored using watchful waiting, and dyspnoea was acutely admitted to the emergency department with stridor. The patient was unresponsive to verbal stimulation. Initial physical examination revealed no signs of symptoms from other organ systems, including the abdomen. Blood investigation showed C-reactive protein 223, white blood cell count 16.7, haemoglobin 7.8 mmol/L. The initial blood gas analysis showed pH 7.15, pCO₂ 10.4, pO₂ 12.2 and lactate 2.6. The mixed acidosis might have been due to the respiratory and metabolic nature of the patient's disease.

After receiving Continuous Positive Airway Pressure (CPAP), the patient became responsive to verbal stimulation. However, the patient still had severe respiratory difficulties and was directly transferred to the intensive care unit to continue CPAP treatment. Suspecting obstructed airways, a chest X-ray was taken. It showed pneumonia, cardiac hypertrophy, a paraesophageal hernia, and a displaced trachea.

The following morning, a Computerized Tomography (CT) scan of the thorax was done in order to clarify the circumstances around the trachea. It confirmed a sizeable paraesophageal hernia pressing onto the trachea, compromising respiratory function. Suspecting the paraesophageal hernia being the reason for the patient's respiratory difficulties, Department of Gastrointestinal Surgery became responsible for further surgical treatment.

Later that day, the patient's condition deteriorated with severe stridor, and an endotracheal tube was inserted.

The next morning, it was decided to operate acutely. In the meantime, the CT scan of the thorax had raised suspicion of ileus, because it showed dilated small bowel segments with air-fluid levels. A thoracoabdominal CT scan was done prior to surgery in order to clarify this. It revealed appendicitis and ileus (Fig. 1) and confirmed the thoracic findings of the earlier CT scan (Fig. 2).

Suspecting an abdominal cause for the patient's symptoms, a diagnostic laparoscopy was performed. It confirmed the suspicion that perforated appendicitis with secondary diffuse peritonitis and ileus had caused high intra-abdominal pressure, resulting in further herniation of the paraesophageal hernia. This made the paraesophageal hernia press onto the trachea, resulting in acute compromised respiratory function. Conversion to explorative laparotomy was necessary due to limited space in the abdominal cavity, which was filled by dilated small bowel segments. Appendectomy and gastropexy ad modum Borema was performed.

Postoperatively, the patient had a 6-day stay in the intensive care unit, to be respiratory stabilized and extubated. The following postoperative course was uneventful. The patient was discharged on the 12th postoperative day and transferred to rehabilitation (Table 1).

3. Discussion

This case reports an acute complication to a paraesophageal hernia requiring emergency surgery, which, to our knowledge, is not previously reported. The hypothesis is that perforated appendicitis caused secondary diffuse peritonitis and ileus. This condition led to increased intra-abdominal pressure. The increased pressure caused a greater amount of stomach to herniate into the thorax, thereby causing an expansion of the existing paraesophageal hernia. This expansion resulted in compression of the trachea, and consequently acute compromised respiratory function.

Download English Version:

<https://daneshyari.com/en/article/5732941>

Download Persian Version:

<https://daneshyari.com/article/5732941>

[Daneshyari.com](https://daneshyari.com)