Accepted Manuscript

Title: Acute intravenous administration of dietary constituent theanine suppresses noxious neuronal transmission of trigeminal spinal nucleus caudalis in rats

Authors: Shiori Takehana, Yoshiko Kubota, Nobuo Uotsu, Kei Yui, Yoshihito Shimazu, Mamoru Takeda

PII: \$0361-9230(16)30354-9

DOI: http://dx.doi.org/doi:10.1016/j.brainresbull.2017.03.004

Reference: BRB 9183

To appear in: Brain Research Bulletin

Received date: 18-10-2016 Revised date: 8-2-2017 Accepted date: 9-3-2017

Please cite this article as: Shiori Takehana, Yoshiko Kubota, Nobuo Uotsu, Kei Yui, Yoshihito Shimazu, Mamoru Takeda, Acute intravenous administration of dietary constituent theanine suppresses noxious neuronal transmission of trigeminal spinal nucleus caudalis in rats, Brain Research Bulletin http://dx.doi.org/10.1016/j.brainresbull.2017.03.004

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

Acute intravenous administration of dietary constituent theanine suppresses noxious neuronal transmission of trigeminal spinal nucleus caudalis in rats

Shiori Takehana ^a, Yoshiko Kubota ^b, Nobuo Uotsu ^b, Kei Yui ^b, Yoshihito Shimazu ^a and Mamoru Takeda ^{a*}

^a Laboratory of Food and Physiological Sciences, Department of Life and Food Sciences, School of Life and Environmental Sciences, Azabu University, 1-17-71, Fuchinobe, Chuo-ku, Sagamihara, Kanagawa, 252-5201, Japan.

FANCL Health Science Research Center, Research Institute, FANCL Corporation, 12-13, Kamishinano, Totsuka-ku, Yokohama, Kanagawa, 244-0806

*Corresponding author: Mamoru Takeda, D.V.M., Ph.D., Professor

Laboratory of Food and Physiological Sciences,

Department of Life and Food Sciences,

School of Life and Environmental Sciences,

Azabu University, 1-17-71, Fuchinobe,

Chuo-ku, Sagamihara, Kanagawa, 252-5201, Japan.

Tel: -81-42-769-1886 Fax:-81-42-769-2212

E-mail: m-takeda@azabu-u.ac.jp

Highlights

- Intravenous theanine injection suppresses nociceptive SpVc WDR neuronal excitability.
- Neuronal firing rates were dose-dependently inhibited by theanine.
- The inhibitory effect of theanine lasted for 5-10 min and was reversible.
- Theanine inhibited the iontophoretic glutamate-induced neuronal firing.
- Theanine may be an effective treatment option for trigeminal pain without side-effects.

Download English Version:

https://daneshyari.com/en/article/5736264

Download Persian Version:

https://daneshyari.com/article/5736264

<u>Daneshyari.com</u>