

## Accepted Manuscript

Title: Amygdala-Prefrontal Cortex Resting-State Functional Connectivity Varies with First Depressive or Manic Episode in Bipolar Disorder

Authors: Shengnan Wei, Haiyang Geng, Xiaowei Jiang, Qian Zhou, Miao Chang, Yifang Zhou, Ke Xu, Yanqing Tang, Fei Wang



PII: S0304-3940(17)30072-1  
DOI: <http://dx.doi.org/doi:10.1016/j.neulet.2017.01.052>  
Reference: NSL 32594

To appear in: *Neuroscience Letters*

Received date: 14-11-2016  
Revised date: 20-1-2017  
Accepted date: 21-1-2017

Please cite this article as: Shengnan Wei, Haiyang Geng, Xiaowei Jiang, Qian Zhou, Miao Chang, Yifang Zhou, Ke Xu, Yanqing Tang, Fei Wang, Amygdala-Prefrontal Cortex Resting-State Functional Connectivity Varies with First Depressive or Manic Episode in Bipolar Disorder, *Neuroscience Letters* <http://dx.doi.org/10.1016/j.neulet.2017.01.052>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Highlights**

This study used functional connectivity techniques to clarify mood state-related features.

The individuals with BD were at their first elevated and depressed mood episode.

To examine the amygdala-PFC neural system was to identify possible state-related neural system patterns at rest in BD.

Download English Version:

<https://daneshyari.com/en/article/5738504>

Download Persian Version:

<https://daneshyari.com/article/5738504>

[Daneshyari.com](https://daneshyari.com)