

Accepted Manuscript

Counter-diffusion biofilms have lower N₂O emissions than co-diffusion biofilms during simultaneous nitrification and denitrification: Insights from depth-profile analysis

Co Thi Kinh, Toshikazu Suenaga, Tomoyuki Hori, Shohei Riya, Masaaki Hosomi, Barth F. Smets, Akihiko Terada

PII: S0043-1354(17)30630-9

DOI: [10.1016/j.watres.2017.07.058](https://doi.org/10.1016/j.watres.2017.07.058)

Reference: WR 13101

To appear in: *Water Research*


Received Date: 3 February 2017

Revised Date: 18 July 2017


Accepted Date: 23 July 2017

Please cite this article as: Kinh, C.T., Suenaga, T., Hori, T., Riya, S., Hosomi, M., Smets, B.F., Terada, A., Counter-diffusion biofilms have lower N₂O emissions than co-diffusion biofilms during simultaneous nitrification and denitrification: Insights from depth-profile analysis, *Water Research* (2017), doi: 10.1016/j.watres.2017.07.058.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Graphical abstract



Download English Version:

<https://daneshyari.com/en/article/5758734>

Download Persian Version:

<https://daneshyari.com/article/5758734>

[Daneshyari.com](https://daneshyari.com)