

# Accepted Manuscript

Effects of decreasing activated carbon particle diameter from 30  $\mu\text{m}$  to 140 nm on equilibrium adsorption capacity

Long Pan, Yuki Nishimura, Hideki Takaesu, Yoshihiko Matsui, Taku Matsushita, Nobutaka Shirasaki



PII: S0043-1354(17)30648-6

DOI: [10.1016/j.watres.2017.07.075](https://doi.org/10.1016/j.watres.2017.07.075)

Reference: WR 13118

To appear in: *Water Research*


Received Date: 29 March 2017

Revised Date: 28 July 2017

Accepted Date: 30 July 2017

Please cite this article as: Pan, L., Nishimura, Y., Takaesu, H., Matsui, Y., Matsushita, T., Shirasaki, N., Effects of decreasing activated carbon particle diameter from 30  $\mu\text{m}$  to 140 nm on equilibrium adsorption capacity, *Water Research* (2017), doi: 10.1016/j.watres.2017.07.075.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Download English Version:

<https://daneshyari.com/en/article/5758741>

Download Persian Version:

<https://daneshyari.com/article/5758741>

[Daneshyari.com](https://daneshyari.com)