Oxalic Acid Produced by *Aspergillus niger* Y-1 Is Effective for Suppression of Bacterial Fruit Blotch of Watermelon Seedlings

Tianyi Gao, Fangmin Hao, Dan Yang, Zhinong Bie, Guoqing Li

PII: S1049-9644(17)30124-X
DOI: http://dx.doi.org/10.1016/j.biocontrol.2017.06.001
Reference: YBCON 3599

To appear in: *Biological Control*

Received Date: 24 January 2017
Revised Date: 4 June 2017
Accepted Date: 6 June 2017

Please cite this article as: Gao, T., Hao, F., Yang, D., Bie, Z., Li, G., Oxalic Acid Produced by *Aspergillus niger* Y-1 Is Effective for Suppression of Bacterial Fruit Blotch of Watermelon Seedlings, *Biological Control* (2017), doi: http://dx.doi.org/10.1016/j.biocontrol.2017.06.001

Oxalic Acid Produced by *Aspergillus niger* Y-1 Is Effective for Suppression of Bacterial Fruit Blotch of Watermelon Seedlings

Tianyi Gao[1], Fangmin Hao[1], Dan Yang[1], Zhinong Bie[2], Guoqing Li[1*]

[1]*State Key Laboratory of Agricultural Microbiology and Key Laboratory of Plant Pathology of Hubei Province*, *Huazhong Agricultural University*, *Wuhan 430070*, *China*; *and* [2]*The Key Lab Horticulture Plant Biology of Ministry of Education, Huazhong Agriculture University, Wuhan 430070, Hubei, China*

[*]Corresponding author: Dr. Guoqing Li, E-mail: guoqingli@mail.hzau.edu.cn