## Accepted Manuscript

Nutritional alterations and damages to stored chickpea in relation with the pest status of Callosobruchus maculatus (Chrysomelidae)



Soumaya Haouel Hamdi, Sourour Abidi, Dorra Sfayhi, Mohamed Zied Dhraief, Moez Amri, Emna Boushih, Mariam Hedjal-Chebheb, Khouja Mouhamed Larbi, Jouda Mediouni Ben Jemâa

PII:	S1226-8615(17)30160-7
DOI:	doi: 10.1016/j.aspen.2017.08.008
Reference:	ASPEN 1032
To appear in:	Journal of Asia-Pacific Entomology
Received date:	13 March 2017
Revised date:	12 July 2017
Accepted date:	16 August 2017

Please cite this article as: Soumaya Haouel Hamdi, Sourour Abidi, Dorra Sfayhi, Mohamed Zied Dhraief, Moez Amri, Emna Boushih, Mariam Hedjal-Chebheb, Khouja Mouhamed Larbi, Jouda Mediouni Ben Jemâa , Nutritional alterations and damages to stored chickpea in relation with the pest status of Callosobruchus maculatus (Chrysomelidae), *Journal of Asia-Pacific Entomology* (2017), doi: 10.1016/j.aspen.2017.08.008

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## **ACCEPTED MANUSCRIPT**

## Nutritional alterations and damages to stored chickpea in relation with the Pest status of *Callosobruchus maculatus* (Chrysomelidae)

Soumaya Haouel Hamdi<sup>1</sup>, Sourour Abidi<sup>2</sup>, Dorra Sfayhi<sup>3</sup>, Mohamed Zied Dhraief<sup>4</sup>, Moez Amri<sup>5</sup>, Emna Boushih<sup>1</sup>, Mariam Hedjal-Chebheb<sup>6</sup>, Khouja Mouhamed Larbi<sup>7</sup>, Jouda Mediouni Ben Jemâa<sup>1,\*</sup>

<sup>1</sup> Laboratory of Biotechnology Applied to Agriculture, National Agricultural Research Institute of Tunisia (INRAT), Rue Hedi Karray, 2080 Ariana, Tunis, University of Carthage, Tunisia.

<sup>2</sup> Laboratory of Animal and Forage Production, National Agricultural Research Institute of Tunisia (INRAT), Rue Hedi Karray, 2080 Ariana, Tunis, University of Carthage, Tunisia.

<sup>3</sup> Field Crop Laboratory, National Agricultural Research Institute of Tunisia (INRAT), Rue Hedi Karray, 2080 Ariana, Tunis, University of Carthage, Tunisia.

<sup>4</sup> National Agricultural Research Institute of Tunisia (INRAT), Rue Hedi Karray, 2080 Ariana, Tunis, University of Carthage, Tunisia.

<sup>5</sup> Biodiversity and Integrated Gene Management Program (BIGM), International Center for Agricultural Research in the Dry Areas (ICARDA), Avenue Mohamed Belarbi Alaoui, BP 6299, Al-Irfane Rabat, Morocco. former researcher at the Field Crop Laboratory, Regional Field Crop Research Center of Beja, Tunisia (CRRGC) and National Institute of Agricultural Research of Tunisia (INRAT).

<sup>6</sup>Faculty of Biological and Agricultural Sciences, University Mouloud Mammeri, Tizi Ouzou, Algeria.

<sup>7</sup> National Institute for Research in Rural Engineering, Water and Forests, INRGREF , University of Carthage, Tunisia. , Tunisia Download English Version:

## https://daneshyari.com/en/article/5763451

Download Persian Version:

https://daneshyari.com/article/5763451

Daneshyari.com