


Accepted Manuscript

Extreme waves in New Zealand waters

Victor A. Godoi , Karin R. Bryan , Scott A. Stephens ,
Richard M. Gorman

PII: S1463-5003(17)30118-X
DOI: [10.1016/j.ocemod.2017.08.004](https://doi.org/10.1016/j.ocemod.2017.08.004)
Reference: OCEMOD 1234



To appear in: *Ocean Modelling*

Received date: 19 December 2016
Revised date: 27 July 2017
Accepted date: 3 August 2017

Please cite this article as: Victor A. Godoi , Karin R. Bryan , Scott A. Stephens ,
Richard M. Gorman , Extreme waves in New Zealand waters, *Ocean Modelling* (2017), doi:
[10.1016/j.ocemod.2017.08.004](https://doi.org/10.1016/j.ocemod.2017.08.004)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights:

- Extreme and mean Hs show similar patterns in most regions around New Zealand (NZ)
- Increasing frequency of extreme waves on the east and south coasts of NZ
- Extreme Hs is likely to be influenced by La Niña-like and SAM effects around NZ
- Extreme Hs has little seasonal variation around the NZ coastline

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/5766345>

Download Persian Version:

<https://daneshyari.com/article/5766345>

[Daneshyari.com](https://daneshyari.com)