

Accepted Manuscript

Statistical Models of Global Langmuir Mixing


Qing Li, Baylor Fox-Kemper, Øyvind Breivik, Adrean Webb

PII: S1463-5003(17)30047-1
DOI: [10.1016/j.ocemod.2017.03.016](https://doi.org/10.1016/j.ocemod.2017.03.016)
Reference: OCEMOD 1195

To appear in: *Ocean Modelling*

Received date: 4 October 2016
Revised date: 20 March 2017
Accepted date: 30 March 2017

Please cite this article as: Qing Li, Baylor Fox-Kemper, Øyvind Breivik, Adrean Webb, Statistical Models of Global Langmuir Mixing, *Ocean Modelling* (2017), doi: [10.1016/j.ocemod.2017.03.016](https://doi.org/10.1016/j.ocemod.2017.03.016)



This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- Two statistical models of global Langmuir mixing are explored and tested.
- Langmuir mixing effects are reproduced with significantly less computational cost.
- The consequences of Stokes depth and misaligned wind and waves are important.

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/5766410>

Download Persian Version:

<https://daneshyari.com/article/5766410>

[Daneshyari.com](https://daneshyari.com)