

Accepted Manuscript

Discrimination of Gluten-Free Oats from Contaminants Using Near Infrared Hyperspectral Imaging Technique

Chyngyz Erkinbaev, Kelly Henderson, Jitendra Paliwal



PII: S0956-7135(17)30223-2
DOI: 10.1016/j.foodcont.2017.04.036
Reference: JFCO 5590
To appear in: *Food Control*
Received Date: 18 January 2017
Revised Date: 15 March 2017
Accepted Date: 23 April 2017

Please cite this article as: Chyngyz Erkinbaev, Kelly Henderson, Jitendra Paliwal, Discrimination of Gluten-Free Oats from Contaminants Using Near Infrared Hyperspectral Imaging Technique, *Food Control* (2017), doi: 10.1016/j.foodcont.2017.04.036

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Research highlights:

Near infrared hyperspectral imaging technique was used to discriminate gluten-free oats.

Two classification methods (PCA and PLSDA) were assessed for accurate grain classification.

Five most significant wavelengths were selected to successfully discriminate oat kernels.

Oats and barley samples (with husk) had similar spectral features resulting in misclassification.

Hyperspectral imaging has potential to be used as a reference tool for quality control in gluten-free oat processing facilities.

Download English Version:

<https://daneshyari.com/en/article/5767243>

Download Persian Version:

<https://daneshyari.com/article/5767243>

[Daneshyari.com](https://daneshyari.com)