



Center problem for trigonometric Liénard systems

Armengol Gasull^a, Jaume Giné^{b,*}, Claudia Valls^c

^a *Departament de Matemàtiques, Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Catalonia, Spain*

^b *Departament de Matemàtica, Inspires Research Centre, Universitat de Lleida, Avda. Jaume II, 69, 25001 Lleida, Catalonia, Spain*

^c *Departamento de Matemática, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais, 1049-001, Lisboa, Portugal*

Received 15 December 2016; revised 10 May 2017

Abstract

We give a complete algebraic characterization of the non-degenerated centers for planar trigonometric Liénard systems. The main tools used in our proof are the classical results of Cherkas on planar analytic Liénard systems and the characterization of some subfields of the quotient field of the ring of trigonometric polynomials. Our results are also applied to some particular subfamilies of planar trigonometric Liénard systems. The results obtained are reminiscent of the ones for planar polynomial Liénard systems but the proofs are different.

© 2017 Elsevier Inc. All rights reserved.

MSC: primary 34C25; secondary 37C10, 37C27

Keywords: Center problem; Trigonometric Liénard equation; Trigonometric polynomial

1. Introduction and main results

The aim of this paper is to characterize the non-degenerated centers for the planar systems associated to the second order trigonometric Liénard differential equations $\ddot{\theta} = g(\theta) + f(\theta)\dot{\theta}$,

* Corresponding author.

E-mail addresses: gasull@mat.uab.cat (A. Gasull), gine@matematica.udl.cat (J. Giné), cvals@math.ist.utl.pt (C. Valls).

<http://dx.doi.org/10.1016/j.jde.2017.05.008>

0022-0396/© 2017 Elsevier Inc. All rights reserved.

where f, g are trigonometric polynomials with real coefficients and the dot denotes the derivative with respect to the time.

The analysis of equations of this form is motivated by a number of problems resulting from pendulum-like equations appearing in the literature. Equations of this form, like $\ddot{\theta} + \sin(\theta) = \varepsilon \dot{\theta} \cos(n\theta)$, or the Josephson equation $\ddot{\theta} + \sin(\theta) = \varepsilon[a - (1 + \gamma \cos(\theta))\dot{\theta}]$, are considered in [9,16,17] or [2,14,18,19], respectively. Also the perturbed whirling pendulum, $\ddot{\theta} = \sin\theta(\cos\theta - \gamma) + \varepsilon(\cos\theta + \alpha)\dot{\theta}$, see [15] falls in this class. Here a, γ, α and ε are real constants and $n \in \mathbb{N}$.

As usual we will write the above second order trigonometric differential equation as the autonomous planar system

$$\begin{cases} \dot{\theta} = y, \\ \dot{y} = g(\theta) + yf(\theta), \end{cases} \quad (1)$$

and we will assume that $f(0) = 0$, $g(0) = 0$, $g'(0) < 0$, where the prime denotes the derivative with respect to θ . These hypotheses on g imply that the origin is either a center or a weak focus. Our main result is:

Theorem 1. *System (1) has a center at the origin if and only if*

- (i) *Either $f = \alpha g$ for some $\alpha \in \mathbb{R}$, or*
- (ii) *There exist a real trigonometric polynomial p and two real polynomials f_1 and g_1 satisfying $p'(0) = 0$, $g_1(p(0))p''(0) < 0$, and such that*

$$f(\theta) = f_1(p(\theta))p'(\theta), \quad g(\theta) = g_1(p(\theta))p'(\theta). \quad (2)$$

The proof of this result is based on the following tools. First, the characterization of the non-degenerated centers for analytic differential equations given by Cherkas in [3], developed in [5] and later improved for polynomial Liénard systems in [4]. Secondly, a version of Luröth Theorem (see [20] or Theorem 5 below) for trigonometric polynomials given in [7,13] and stated as Theorem 7. We will recall these results in Section 2.

From condition (2), by introducing the functions

$$F(\theta) = \int_0^\theta f(s) ds, \quad G(\theta) = \int_0^\theta g(s) ds, \quad (3)$$

it follows that, for trigonometric Liénard systems having a center of type (ii),

$$F(\theta) = F_1(p(\theta)), \quad G(\theta) = G_1(p(\theta)), \quad (4)$$

for some polynomial functions F_1 and G_1 , with $p'(0) = 0$, $G_1'(p(0))p''(0) < 0$. This equivalent expression is commonly used to characterize the centers of polynomial Liénard systems, see [4].

We want to remark that the centers of item (i) in the theorem have explicit global first integrals,

$$H(\theta, y) = \begin{cases} y^2 - 2G(\theta), & \text{when } \alpha = 0, \\ (1 + \alpha y) \exp(\alpha^2 G(\theta) - \alpha y), & \text{when } \alpha \neq 0. \end{cases} \quad (5)$$

Download English Version:

<https://daneshyari.com/en/article/5774121>

Download Persian Version:

<https://daneshyari.com/article/5774121>

[Daneshyari.com](https://daneshyari.com)