

RESEARCH

Open Access



# Periodic orbits to Kaplan-Yorke like differential delay equations with two lags of ratio $(2k - 1)/2$

Lin Li<sup>1</sup>, Chunyan Xue<sup>1\*</sup> and Weigao Ge<sup>2</sup>

\*Correspondence:  
xue\_chunyan@126.com  
<sup>1</sup>School of Science, Beijing  
Information Science and  
Technology University, Beijing,  
100192, P.R. China  
Full list of author information is  
available at the end of the article

## Abstract

In this paper, we study the periodic solutions to a type of differential delay equations with two lags of ratio  $(2k - 1)/2$  in the form

$$x'(t) = -f(x(t-2)) - f(x(t-(2k-1))), \quad k \geq 2.$$

The  $4k$ -periodic solutions are obtained by using the variational method and the method of Kaplan-Yorke coupling system. This is a new type of differential-delay equations compared with all previous researches since the ratio of two lags is not an integer. Three functionals are constructed for a discussion on critical points. An example is given to demonstrate our main results.

**MSC:** 34B10; 34B15

**Keywords:** differential delay equation; periodic solutions; critical point theory; variational method

## 1 Introduction

The differential delay equations have useful applications in various fields such as age-structured population growth, control theory, and any models involving responses with nonzero delays [1–5].

Given  $f \in C^0(\mathbb{R}^+, \mathbb{R}^-)$  with  $f(-x) = -f(x)$ ,  $xf(x) > 0$ ,  $x \neq 0$ . Kaplan and Yorke [6] studied the existence of 4-periodic and 6-periodic solutions to the differential delay equations

$$x'(t) = -f(x(t-1)) \tag{1.1}$$

and

$$x'(t) = -f(x(t-1)) - f(x(t-2)), \tag{1.2}$$

respectively. The method they applied is transforming the two equations into adequate ordinary differential equations by regarding the retarded functions  $x(t-1)$  and  $x(t-2)$  as independent variables. They guessed that the existence of  $2(n+1)$ -periodic solution to the

equation

$$x'(t) = - \sum_{i=1}^n f(x(t-i)) \tag{1.3}$$

could be studied under the restriction

$$x(t - (n + 1)) = -x(t),$$

which was proved by Nussbaum [7] in 1978 by use of a fixed point theorem on cones.

After then, a lot of papers [8–21] discussed the existence and multiplicity of  $2(n + 1)$ -periodic solutions to equation (1.3) and its extension

$$x'(t) = - \sum_{i=1}^n \text{grad } F(x(t-i)), \tag{1.4}$$

where  $F \in C^1(\mathbb{R}^N, \mathbb{R}), F(-x) = F(x), F(0) = 0$ .

Recently, Zhang and Ge [22] studied the multiplicity of  $2n$ -periodic solutions to a type of differential delay equations of the form

$$x'(t) = -f(x(t-1)) - f(x(t-n)), \quad n \geq 2, \tag{1.5}$$

and obtained new results.

In this paper, we study the periodic orbits to a type of differential delay equations with two lags of ratio  $(2k - 1)/2$  in the form

$$x'(t) = -f(x(t-2)) - f(x(t-(2k-1))), \quad k \geq 2, \tag{1.6}$$

which is different from (1.3) and can be regarded as a new extension of (1.2). The method applied in this paper is the variational approach in the critical point theory [23, 24].

Since the equation

$$x'(t) = -f(x(t-2)) - f(x(t-2k)), \quad k \geq 2,$$

can be changed into the form of equation (1.5) by the transformation

$$t = 2s, \quad x(t) = x(2s) = y(s), \quad \widehat{f}(y) = 2f(x),$$

this paper completes the research of the equations in the form

$$x'(t) = -f(x(t-2)) - f(x(t-n)), \quad n \geq 3. \tag{1.7}$$

In fact, it follows from

$$y'(s) = 2x'(t) = -2f(x(t-2)) - 2f(x(t-2k)) = -2f(x(2(s-1))) - 2f(x(2(s-k)))$$

Download English Version:

<https://daneshyari.com/en/article/5775268>

Download Persian Version:

<https://daneshyari.com/article/5775268>

[Daneshyari.com](https://daneshyari.com)