


Accepted Manuscript

Effects of active fault types on earthquake-induced deep-seated landslides: A study of historical cases in Japan

Chi-Wen Chen, Tomoyuki Iida, Ryuji Yamada

PII: S0169-555X(16)30436-6
DOI: doi: [10.1016/j.geomorph.2017.07.030](https://doi.org/10.1016/j.geomorph.2017.07.030)
Reference: GEOMOR 6093
To appear in: *Geomorphology*
Received date: 3 June 2016
Revised date: 31 July 2017
Accepted date: 31 July 2017



Please cite this article as: Chi-Wen Chen, Tomoyuki Iida, Ryuji Yamada , Effects of active fault types on earthquake-induced deep-seated landslides: A study of historical cases in Japan, *Geomorphology* (2017), doi: [10.1016/j.geomorph.2017.07.030](https://doi.org/10.1016/j.geomorph.2017.07.030)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Effects of active fault types on earthquake-induced deep-seated
landslides: a study of historical cases in Japan**

Chi-Wen Chen^{a, b, c, *}, Tomoyuki Iida^{a, b}, Ryuji Yamada^a

^aNational Research Institute for Earth Science and Disaster Resilience, Tsukuba,
Ibaraki 305-0006, Japan

^bCenter for Spatial Information Science, University of Tokyo, 5-1-5 Kashiwanoha
Kashiwa, Chiba 277-8568, Japan

^cNational Science and Technology Center for Disaster Reduction, 9F., No. 200, Sec. 3,
Beixin Road, Xindian District, New Taipei City, Taiwan

Download English Version:

<https://daneshyari.com/en/article/5780785>

Download Persian Version:

<https://daneshyari.com/article/5780785>

[Daneshyari.com](https://daneshyari.com)