


Accepted Manuscript

Deep-seated landslides affecting monoclinal flysch morphostructure: Evaluation of LiDAR-derived topography of the highest range of the Czech Carpathians

Michal Břežný, Tomáš Pánek



PII: S0169-555X(16)31160-6
DOI: doi: [10.1016/j.geomorph.2017.02.007](https://doi.org/10.1016/j.geomorph.2017.02.007)
Reference: GEOMOR 5923
To appear in: *Geomorphology*
Received date: 7 December 2016
Revised date: 2 February 2017
Accepted date: 17 February 2017

Please cite this article as: Michal Břežný, Tomáš Pánek , Deep-seated landslides affecting monoclinal flysch morphostructure: Evaluation of LiDAR-derived topography of the highest range of the Czech Carpathians. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Geomor(2017), doi: [10.1016/j.geomorph.2017.02.007](https://doi.org/10.1016/j.geomorph.2017.02.007)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Deep-seated landslides affecting monoclinial flysch morphostructure:
evaluation of LiDAR-derived topography of the highest range of the Czech
Carpathians**

Michal Břežný^{1*}, Tomáš Pánek¹

1. Department of Physical Geography and Geoecology, Faculty of Science, University of Ostrava, Chittussiho 10, 710 00 Ostrava, Czech Republic

*Corresponding author. Tel. : +420 774 918 954

E-mail: michal.brezny@osu.cz (M. Břežný)

Download English Version:

<https://daneshyari.com/en/article/5781030>

Download Persian Version:

<https://daneshyari.com/article/5781030>

[Daneshyari.com](https://daneshyari.com)