

## Accepted Manuscript


Desiccation cracks in siliciclastic deposits: Microbial mat-related compared to abiotic sedimentary origin

Olga Kovalchuk, George W. Owttrim, Kurt O. Konhauser, Murray K. Gingras

PII: S0037-0738(16)30279-2  
DOI: doi:[10.1016/j.sedgeo.2016.11.002](https://doi.org/10.1016/j.sedgeo.2016.11.002)  
Reference: SEDGEO 5129

To appear in: *Sedimentary Geology*

Received date: 20 July 2016  
Revised date: 4 November 2016  
Accepted date: 6 November 2016



Please cite this article as: Kovalchuk, Olga, Owttrim, George W., Konhauser, Kurt O., Gingras, Murray K., Desiccation cracks in siliciclastic deposits: Microbial mat-related compared to abiotic sedimentary origin, *Sedimentary Geology* (2016), doi:[10.1016/j.sedgeo.2016.11.002](https://doi.org/10.1016/j.sedgeo.2016.11.002)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# Desiccation cracks in siliciclastic deposits: Microbial mat-related compared to abiotic sedimentary origin

Olga Kovalchuk<sup>a,\*</sup>, George W. Owttrim<sup>b</sup>, Kurt O. Konhauser<sup>a</sup>, Murray K. Gingras<sup>a</sup>

<sup>a</sup> Department of Earth and Atmospheric Sciences, University of Alberta, Edmonton, Alberta, Canada, T6G 2E3

<sup>b</sup> Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada, T6G 2E9

\* Corresponding author. E-mail address: okovalch@ualberta.ca. Fax: 1 (780) 492-2030.

## Abstract

Siliciclastic sediment colonized by microbial mats yield a set of distinct sedimentary fabrics that are collectively called “mat-related structures (MRS)”. In the rock record, versatile cracks are observed in biostabilized strata, but the mechanisms responsible for their formation remain debated. Microbially stabilized sediments produce desiccation cracks that serve as modern analogues for fossil microbial cracks. However, since both microbial mat shrinkage and clay shrinkage may contribute to the formation of these desiccation cracks, it is difficult to isolate the influence of the microbial mat on the resulting crack formation, distribution and morphology. To address this issue, we conducted a series of desiccation experiments that determine differences between microbially influenced desiccation cracks (i.e. biotic) and those formed in identical, but sterilized (i.e. abiotic) siliciclastic sediment. Three sediment mixtures were used: (1) very fine-sized sand, (2) mixed (ungraded) silt/clay, and (3) normally graded silt/clay. In all of the experiments, the water-rich microbial mat contracted substantially while drying, producing isolated pockets of shallow, but wide cracks, the distribution of which was controlled by heterogeneities in the mat structure and thickness variations of the mat. In the sand-rich substratum, the microbial mat was the only crack-forming mechanism, while in the clay-rich substrata (experiments

Download English Version:

<https://daneshyari.com/en/article/5781328>

Download Persian Version:

<https://daneshyari.com/article/5781328>

[Daneshyari.com](https://daneshyari.com)