Accepted Manuscript

Depositional ages and provenance of the Neoproterozoic Damara Supergroup (northwest Namibia): Implications for the Angola-Congo and Kalahari cratons connection

Débora B. Nascimento, Renata S. Schmitt, André Ribeiro, Rudolph A.J. Trouw, Cees W. Passchier, Miguel A.S. Basei

PII: S1342-937X(16)30425-7 DOI: doi:10.1016/j.gr.2017.09.006

Reference: GR 1865

To appear in:

Received date: 25 November 2016

Revised date: 4 July 2017

Accepted date: 24 September 2017

Please cite this article as: Débora B. Nascimento, Renata S. Schmitt, André Ribeiro, Rudolph A.J. Trouw, Cees W. Passchier, Miguel A.S. Basei, Depositional ages and provenance of the Neoproterozoic Damara Supergroup (northwest Namibia): Implications for the Angola-Congo and Kalahari cratons connection. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Gr(2017), doi:10.1016/j.gr.2017.09.006

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT

Depositional ages and provenance of the Neoproterozoic Damara Supergroup (northwest Namibia): implications for the Angola-Congo and Kalahari cratons connection

Débora B. Nascimento^{a,b,*}, Renata S. Schmitt^b, André Ribeiro^b, Rudolph A.J. Trouw^b, Cees W. Passchier^c, Miguel A.S. Basei^d

- ^a Programa de Pós–graduação em Geologia, Universidade Federal do Rio de Janeiro (PPGL/UFRJ), CEP 21941-916, Rio de Janeiro, Brazil
- ^b Instituto de Geociências, Universidade Federal do Rio de Janeiro (IGEO/UFRJ), CEP 21941-916, Rio de Janeiro, Brazil
- ^c Institut für Geowissenschaften, Johannes Gutenberg-Universität Mainz, 55099 Mainz, Germany
- ^d Instituto de Geociências, Universidade de São Paulo, CEP 05508-080, São Paulo, Brazil
- *Corresponding author: e-mail: debora@geologia.ufrj.br; tel: +55 21 39389463, fax: +55 21 39389465.

Abstract

The Damara Orogen is composed of the Damara, Kaoko and Gariep belts developed during the Neoproterozoic Pan-African Orogeny. The Damara Belt contains Neoproterozoic siliciclastic and carbonate successions of the Damara Supergroup that record rift to proto-ocean depositional phases during the Rodinia

Download English Version:

https://daneshyari.com/en/article/5785207

Download Persian Version:

https://daneshyari.com/article/5785207

<u>Daneshyari.com</u>