

# Accepted Manuscript

The charophytes-rich Barremian-Albian record in Jebel Ksaïra (Central Tunisia)

Amal Salmouna, Isabelle Cojan, Jamel Tourir



PII: S1464-343X(17)30117-6

DOI: 10.1016/j.jafrearsci.2017.03.011

Reference: AES 2847

To appear in: *Journal of African Earth Sciences*

Received Date: 05 February 2016

Revised Date: 07 March 2017

Accepted Date: 13 March 2017

Please cite this article as: Amal Salmouna, Isabelle Cojan, Jamel Tourir, The charophytes-rich Barremian-Albian record in Jebel Ksaïra (Central Tunisia), *Journal of African Earth Sciences* (2017), doi: 10.1016/j.jafrearsci.2017.03.011

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Highlights

- The sedimentary environment of the Kebar Formation in Jebel Ksaïra locality:
  - oscillated between a continental domain and a marginal marine to infratidal domain.
  - was cyclic, composed by continental deposition and marginal marine to infratidal deposition.
- Dated by charophytes, the Kebar Formation in Jebel Ksaïra is attributed to Late Barremian-Early Albian.
- The Kebar Formation should have been deposited in Jebel Ksaïra earlier than in Jebel Kebar.

Download English Version:

<https://daneshyari.com/en/article/5785463>

Download Persian Version:

<https://daneshyari.com/article/5785463>

[Daneshyari.com](https://daneshyari.com)