


Accepted Manuscript

Detection and correlation of tephra-derived smectite-rich shear zones by analyzing glass melt inclusions in mineral grains

Eiichi Ishii, Akira Furusawa



PII: S0013-7952(16)30562-2
DOI: doi: [10.1016/j.enggeo.2017.08.010](https://doi.org/10.1016/j.enggeo.2017.08.010)
Reference: ENGEO 4619
To appear in: *Engineering Geology*
Received date: 27 October 2016
Revised date: 29 April 2017
Accepted date: 8 August 2017

Please cite this article as: Eiichi Ishii, Akira Furusawa , Detection and correlation of tephra-derived smectite-rich shear zones by analyzing glass melt inclusions in mineral grains. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Enggeo(2017), doi: [10.1016/j.enggeo.2017.08.010](https://doi.org/10.1016/j.enggeo.2017.08.010)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Detection and correlation of tephra-derived smectite-rich shear zones by analyzing glass melt inclusions in mineral grains

Eiichi Ishii^{1,*} and Akira Furusawa²

¹ *Horonobe Underground Research Laboratory, Japan Atomic Energy Agency, Horonobe-cho, Japan*

² *Furusawa Geological Survey Corporation, Okazaki, Japan*

*Corresponding author. E-mail address: ishii.eiichi@jaea.go.jp

Download English Version:

<https://daneshyari.com/en/article/5787516>

Download Persian Version:

<https://daneshyari.com/article/5787516>

[Daneshyari.com](https://daneshyari.com)