

# Dengue: what it is and why there is more

Yuan Li · Shuyu Wu

Received: 20 January 2015 / Accepted: 27 January 2015  
© Science China Press and Springer-Verlag Berlin Heidelberg 2015

**Abstract** In 2014, China experienced the worst outbreak of dengue fever in the last decade with over 40,000 dengue cases including six deaths by the end of October. As one of the “neglected” tropical diseases, dengue is affecting substantially increasing number of people and proportion of global population due to factors including globalization, human settlement, and possibly climate change. Here, the authors summarized the most recent data about dengue outbreaks in China and reviewed the global trend of dengue epidemiology. Future directions for dengue surveillance, control and prevention are also introduced.

**Keywords** Dengue · Vector control · One Health · China

In 2014, China experienced the worst outbreak of dengue fever (DF) in the last decade (Fig. 1a, b). A total of 44,591 dengue cases were reported to the national notifiable infectious diseases reporting system by the end of October including six deaths, and 99 % of cases were reported in south China provinces, such as Guangdong, Fujian, Yunnan and Zhuang Autonomous Region [1]. On November 3, the U.S. Centers for Disease Control and Prevention (CDC) issued a travel notice for Guangdong province, China because of the dengue outbreak [2].


## 1 The global threat of dengue

Dengue fever (DF), caused by any one of the four closely related dengue viruses (DENV 1–4), is a viral illness transmitted by *Aedes* mosquitoes. Originating in monkeys, dengue viruses were transmitted to humans between 100–800 years ago, and dengue remained a relatively minor and geographically restricted disease until the 1950s when epidemics occurred in the Philippines and Thailand [3, 4]. Dengue has emerged as a worldwide problem, with a 30-fold increase in incidence globally in the past 50 years (Fig. 2) [4]. Today, dengue is regarded as the most prevalent and rapidly spreading mosquito-borne viral disease of humans [4]. It is found in at least 100 tropical and subtropical countries, including countries in Africa, Southeast Asia, the western Pacific, the Americas, the Caribbean, and the eastern Mediterranean regions [3, 5]. More than one-third of the world’s population lives in areas at risk of infection with dengue virus [3], and a recent estimate using cartographic approaches in 2013 showed that 390 million people have dengue virus infections per year, of which 96 million cases manifest symptoms worldwide [6].


Dengue is a complex disease with a wide spectrum of clinical presentation. Classic DF is characterized by acute onset of high fever 3–14 days after the bite of an infected mosquito, and the high fever can persist for 2–7 days accompanied by frontal headache, retro-orbital pain, myalgias, arthralgias, rash, and low white blood cell count [7, 8]. Due to the severe muscle, joint and bone pains, dengue is also known as “break bone fever”. Most dengue patients recover after a few days, but some DF patients develop dengue hemorrhagic fever (DHF) or dengue shock syndrome (DSS), which are severe and sometimes fatal forms of the disease [7]. About three quarters of dengue infections can be asymptomatic, especially in children and those

---

Y. Li (✉) · S. Wu  
Global Disease Detection Program, United States Centers for  
Disease Control and Prevention, Beijing 100600, China  
e-mail: liy@cn.cdc.gov



**Fig. 1** Number of reported dengue fever (DF) cases in China's mainland from 2004 to 2013 (a) and from January to October 2014 (b). Data source: Notifiable infectious disease reporting summary, China CDC [1]



**Fig. 2** Average number of dengue and severe dengue (DHF) cases reported to WHO from 1955 to 2007 [4]

with previous dengue infection, but they can still infect mosquitoes [3, 7]. However, secondary dengue infection is one of the two factors thought to be associated with increased risk of disease severity; the other factor is infection by particularly virulent strains of dengue viruses [8].

At present, no vaccine exists to prevent dengue infection, nor are there specific medications to treat the disease. The most effective protective measures are those that prevent mosquito bites, namely vector control [3]. The main arthropod vector for dengue virus transmission is *Aedes aegypti*, and a second, less effective vector is *Aedes*

*albopictus* [8]. Control of dengue vectors has mainly been addressed by community-based source reduction: The elimination of containers that are favorable sites for oviposition and development of the aquatic stages [4]. For individuals, preventive measures include sleeping under a mosquito bed net, eliminating mosquito indoors and wearing repellent [3].

The World Health Organization (WHO) described dengue as a “neglected” tropical disease (NTD) in its Dengue Prevention and Control Strategy 2012–2020, due to the lack of global coordination efforts, research, and political will [4]. The contemporary worldwide distribution of the risk of dengue virus infection and its public health burden is poorly known, and the greatest uncertainties in national burden estimates are in India, Indonesia, Brazil, and China [6]. The economic burden of dengue on both governments and individuals is high. Based on a systematic review analysis, the estimated cost of dengue disease in the Americas including North America, Central America and Mexico, the Andean region, Brazil, the Southern cone, and the Caribbean region is 2.1 billion USD per year on average, excluding cost on vector control [9]. Another 12-country economic analysis also showed high dengue burden in Southeast Asia, with an annual average of 950 million USD resulted from 2.9 million dengue episodes and 5,906 deaths [10]. Information on economic burden of dengue in China, however, is not available.

## 2 Dengue in China

NTDs including dengue are still widespread and create a substantial burden in China [11]. Infectious diseases in China are divided into three categories (A, B, and C) according to their transmission modes, speed, and hazard to humans; DF, together with Severe Acute Respiratory Syndrome (SARS), H1N1 influenza, rabies, tuberculosis etc., belong to infectious diseases category B. In 2005, China established a national monitoring program for DF in humans and vector monitoring in *Aedes* mosquitoes, focusing on five provinces (Guangdong, Fujian, Yunnan, Guangxi Autonomous Region, and Hainan) with 16 national monitoring sites. Other efforts that China has made to prevent and control dengue include control of mosquito vector density, enhanced surveillance for imported cases, improvement of diagnosis and treatment, enhancing the cooperation among the related different public health departments and neighboring countries, and engaging communities to combat the epidemic [11, 12]. In response to the dengue outbreak in 2014, four technical guidelines [13], including surveillance of DF cases, laboratory tests and diagnosis, surveillance of dengue vectors, and control of dengue vectors, were developed and issued by China

Download English Version:

<https://daneshyari.com/en/article/5789266>

Download Persian Version:

<https://daneshyari.com/article/5789266>

[Daneshyari.com](https://daneshyari.com)