




ELSEVIER

Contents lists available at ScienceDirect

Livestock Science

journal homepage: www.elsevier.com/locate/livsci

Short communication

Association between hematological status at weaning and weight gain post-weaning in piglets



Sheeva Bhattarai*, Jens Peter Nielsen

HERD – Centre for Herd-oriented Education, Research and Development, Department of Large Animal Sciences, University of Copenhagen, Groennegaardsvej 2, 1870 Frederiksberg C, Denmark

ARTICLE INFO

Article history:

Received 13 April 2015

Received in revised form

19 October 2015

Accepted 21 October 2015

Keywords:

Anemia

Growth

Hemoglobin

Iron

Pig

ABSTRACT

The objective of the study was to determine the associations between hematological parameters at weaning and post-weaning weight gain in piglets. Five well-managed conventional herds of average herd size of 1000 sows were selected. All herds provided a single injectable iron (either i.m. or s.c, 200 mg) supplementation from different commercial products at 3–4 days of life. Within each herd, litters belonging to a weekly farrowing batch close to weaning were identified and among them 20 litters were chosen randomly after exclusion of nursing sows. Within each litter, a random piglet was selected. EDTA and non-stabilized blood sample was taken from the anterior Vena cava of each piglet 1–3 days before weaning (average weaning age of 25.7 days) and analyzed by complete hematology including serum iron and total iron binding capacity (TIBC). The associations between each measured hematological parameter and the average daily gain (ADG) of piglets in a three-week period were tested using PROC MIXED procedure in SAS. A total of 99 piglets were included in the study but 3 were lost during follow up. The average hemoglobin concentration of piglets was $121.5 \pm 15 \text{ g l}^{-1}$ and the ADG was $202.9 \pm 78.4 \text{ g day}^{-1}$. A positive association between hemoglobin and ADG in the three-week period ($p=0.0003$) was observed. Also, red blood cells ($p < 0.0001$) and hematocrit ($p=0.0005$) had positive association with ADG while red blood cell distribution width ($p=0.05$), hemoglobin distribution width ($p=0.04$), and reticulocyte red cell distribution width ($p=0.01$) had a negative association with ADG. Hence the results indicate that improved hematological status at weaning is positively correlated to weight gain post-weaning. An increase in 10 g hemoglobin/l blood corresponded to a weight gain improvement of 17.2 g daily weight gain in the 3 weeks post-weaning period. The strategies for improving the hematological values of piglets at weaning need to be further investigated.

© 2015 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Piglets are prone to iron deficiency anemia for several reasons including, very low iron stores at birth, inadequate supply from sow milk, and rapid growth rate (Venn et al., 1947; Rincker et al., 2004). Therefore, it is a standard procedure to iron supplement piglets in the first days of life and in some herds also later during the suckling period. However, it is unknown whether optimal hematological values to promote rapid growth of piglets are achieved under production conditions.

Several studies have focussed on the level of hemoglobin and the cut-off values for anemia in piglets (Furugouri, 1975; Egeli and Framstad, 1998). However, even though hemoglobin (Hb) values above anemia cut-off is reached, there may still be a positive production effect of somewhat higher values. Aukett et al. (1986)

found an increased rate of weight gain in children treated with iron compared to untreated children. Gentry et al. (1997) have demonstrated that piglets with greater Hb levels at weaning had greater average daily gain post-weaning. The effect of other hematological parameters on the growth rate of pigs has not been investigated yet.

Some studies have suggested that Hb is not a sensitive indicator of early fall in iron status (Cook, 2005; Svoboda et al., 2008) and several pig and human studies suggest that mean corpuscular volume, mean corpuscular hemoglobin, red cell distribution width, serum iron, total iron binding capacity, transferrin saturation, serum ferritin and reticulocyte parameters are earlier indicators of iron deficiency (Lipschitz et al., 1974; Furugouri et al., 1982; Smith et al., 1984; McClure et al., 1985; Macdougall et al., 1992; Andrews et al., 1994; Ilić et al., 2006; Mast et al., 2008; Svoboda et al., 2008). We hypothesize that Hb and hematological parameters influence the growth rate of piglets and therefore, the aim of this study was to investigate the associations between hematological parameters at weaning and post-weaning weight gain in piglets.

* Corresponding author.

E-mail address: sheeva@sund.ku.dk (S. Bhattarai).

2. Materials and methods

This study was conducted in accordance with the guidelines of the Danish Ministry of Justice Act No. 382 (June 10, 1987) and Acts 333 (May 19, 1990), 726 (September 9, 1993), and 1016 (December 12, 2001) with respect to animal experimentation and care of animals under study.

Five high-performing conventional farrow-finish sow herds were selected for the study (Table 1). Within each herd, 20 litters ready to weaning within 1–3 days and belonging to the same farrowing batch were selected by convenience sampling. The average weaning age of piglets in each herd is presented in Table 2. In each litter, a random piglet was selected as the sixth clinically healthy individual when counting snouts from the pen side of the observer. In total, 99 piglets were studied. Blood was collected from each selected piglet and subjected to complete hematological analysis including serum iron. The procedures of herd selection, sow and piglet selection and method of blood sampling have previously been described in details in Bhattarai and Nielsen (2015). The present study constitutes a follow-up study for a fraction of the piglets described in Bhattarai and Nielsen (2015).

The following recordings were made within each litter: Farrowing date, litter size as counted at the day of visit, sex of the selected piglet, and parity of the sow. The selected piglets were weighed individually and examined for obvious signs of anemia (pale skin) or clinical signs of disease at the time of blood collection. The piglets were ear tagged with individual identification numbers for follow up.

Three weeks after weaning, body weight of all the piglets was recorded again. The average daily gain (ADG) was calculated as the difference between the weaning weight and the 3-week post-weaning weight, divided by the number of days between weighing.

2.1. Hematology

The EDTA stabilized blood samples were analyzed for hematological indices; erythrocyte count (RBC), leucocyte count (total leucocyte count and differential leucocyte count), platelets, mean platelet volume (MPV), red blood cell distribution width (RDW), hemoglobin concentration (Hb), hemoglobin distribution width (HDW), hematocrit (HCT), mean cell volume (MCV), mean corpuscular hemoglobin (MCH), and mean cell hemoglobin concentration (MCHC). Reticulocyte indices were also analyzed which included reticulocyte count (absolute and relative), reticulocyte

Table 1
Descriptive data of the herds that participated in the study.

	Herd 1	Herd 2	Herd 3	Herd 4	Herd 5
Herd size	1101	1100	1155	1201	940
Weaning age (day)	26.3	24.2	27.8	24.8	25.7
Total born alive/litter (piglets)	15.7	15.6	15.7	14.5	14.3
Age at iron injection (day)	3–4	3–4	4	4	3
Iron brand name ¹	Solofer ^a	Ursoferran ^b	Ursoferran ^b	Hyofer ^c	Hyofer ^c
Dose/Route of iron ²	1 ml/i.m.	1.1 ml/i.m.	1 ml/i.m.	1 ml/s.c.	1.5 ml/i.m.
Creep feed start	5 days	7 days	10 days	6 days	2 wks

¹ Each injectable iron product contains 200 mg iron dextran per ml.

² i.m. = Intramuscular; s.c. = Subcutaneous.

^a Pharmacosmos A/S, Holbæk, Denmark.

^b Serumwerk Bernburg AG, Bernburg, Germany.

^c Salfarm Danmark A/S, Kolding, Denmark.

Table 2
Description of piglet growth performance by herd^a.

Herd	No. of piglets studied	Age at weaning (days)	Weight at weaning (kg)	Weight post-weaning (kg)	ADG ^b (g/day)
1	20	26.2 ± 0.6	6.9 ± 1.2	11.3 ± 1.9	209.3 ± 56.2 ^a
2	18	24.2 ± 1.5	6.0 ± 1.8	10.4 ± 2.8	210 ± 66.0 ^a
3	17	27.8 ± 3.4	6.1 ± 1.2	11.1 ± 2.5	232.7 ± 79.4 ^a
4	19	24.8 ± 1.5	5.4 ± 1.0	10.6 ± 2.5	245.5 ± 86.1 ^a
5	19	25.7 ± 0.8	6.5 ± 1.1	9.4 ± 1.4	118.8 ± 31.0 ^b
Mean		25.7 ± 2.2	6.2 ± 1.3	10.6 ± 2.3	202.9 ± 78.4

^aDifferent superscripts within ADG column indicate significant difference ($p < 0.05$).

^a Values are presented as mean ± standard deviations.

^b ADG = Average daily gain. All the piglets were weighed 3 weeks after weaning.

hemoglobin content (Chr), mean reticulocyte corpuscular hemoglobin concentration (CHCMr), reticulocyte cellular volume (MCVr), reticulocyte red cell distribution width (RDWr), and reticulocyte hemoglobin distribution width (HDWr). The serum samples were analyzed for serum iron and total iron binding capacity (TIBC). Transferrin saturation (TfS) was calculated using the formula: TfS (%) = Serum iron / TIBC × 100.

Complete hematology was tested using Advia 2120i Hematology System (Siemens healthcare diagnostics Inc. Tarrytown, NY 10591, United States) while serum iron and TIBC were tested using Advia 1800 Chemistry System (Siemens healthcare diagnostics Inc. Tarrytown, NY 10591, United States) at the Central laboratory, Department of veterinary clinical and animal sciences, University of Copenhagen. Method validations were according to protocols of the manufacturer (Siemens healthcare diagnostics, 2008).

2.2. Statistical analysis

Data analysis was performed using SAS (SAS 9.3, SAS institute Inc, Cary, North Carolina). The associations of each of the measured hematological parameters on the ADG of piglets were tested individually using linear mixed model with PROC MIXED procedure in separate models and the estimates were calculated using SOLUTION option. Farm was considered as a random variable during each analysis with weight at weaning, sex, parity of sow, and litter size as other explanatory variables. The difference in ADG among farms was determined using general linear model with PROC GLM. Pearson's correlation coefficients between hematological parameters were calculated using PROC CORR procedure. Statistical significance was set at $p < 0.05$ for all the tests.

3. Results

Altogether 99 piglets were included in the study. Serum sample was missing from six piglets. The results from 15 EDTA blood samples were not included in the analysis (EDTA blood either not taken/analyzed or clotted). Three piglets were lost during follow up. The average daily gain of the piglets was 202.9 ± 78.4 g day⁻¹ (Table 2) and the mean Hb concentration was 121.5 ± 15 g l⁻¹ (Table 3). The mean values of selected hematological parameters (those showing association with growth rates) are shown in Table 3. Correlations among Hb, RBC, HCT, RDW, HDW and RDWr are only reported as these parameters showed association with growth rates (Table 5).

The associations between individual hematological parameters and ADG of piglets are presented in Table 4. Only statistically significant parameters are presented. ADG post-weaning had a significant association with Hb (Figs. 1 and 2), RBC, HCT, RDW,

Download English Version:

<https://daneshyari.com/en/article/5789982>

Download Persian Version:

<https://daneshyari.com/article/5789982>

[Daneshyari.com](https://daneshyari.com)