

## Vascular damage: A persisting pathology common to Alzheimer's disease and traumatic brain injury



Max Franzblau, Chiara Gonzales-Portillo, Gabriel S. Gonzales-Portillo, Theo Diamandis, Mia C. Borlongan, Naoki Tajiri\*, Cesar V. Borlongan\*

Center of Excellence for Aging and Brain Repair, Department of Neurosurgery and Brain Repair, University of South Florida Morsani College of Medicine, 3515 E. Fletcher Avenue, Tampa, FL 33613, USA

### ARTICLE INFO

#### Article history:

Received 8 August 2013

Accepted 8 September 2013

### ABSTRACT

Alzheimer's disease (AD) and traumatic brain injury (TBI) are both significant clinical problems characterized by debilitating symptoms with limited available treatments. Interestingly, both neurological diseases are characterized by neurovascular damage. This impaired brain vasculature correlates with the onset of dementia, a symptom associated with hippocampal degeneration seen in both diseases. We posit that vascular damage is a major pathological link between TBI and AD, in that TBI victims are predisposed to AD symptoms due to altered brain vasculature; vice versa, the progression of AD pathology may be accelerated by TBI especially when the brain insult worsens hippocampal degeneration. Our hypothesis is supported by recent data reporting expedited AD pathology in presymptomatic transgenic AD mice subjected to TBI. If our hypothesis is correct, treatments targeted at repairing the vasculature may prove effective at treating both diseases and preventing the evolution of AD symptoms in TBI victims.

© 2013 Elsevier Ltd. All rights reserved.

### Alzheimer's disease and vascular damage

Each year, Alzheimer's disease (AD) affects over five million people in the United States. In most cases, symptoms emerge after the age of sixty, and progress on a spectrum of three stages. Memory loss and poor judgment are among the most prominent early signs and symptoms. AD is a neurodegenerative disorder [1] primarily tied to the hippocampus region of the brain with patients exhibiting an inability to communicate and a lack of control of bowel and bladder. As the disease progresses, brain tissue in the surrounding hippocampal areas undergoes similar neurodegeneration, characterized by aberrant tau and amyloid protein deposits [1]. AD is a debilitating, irreversible brain condition that eventually leads to death.

AD's major pathological feature is the degeneration and loss of cholinergic neurons and synapses [2], which compromises the brain. This neuronal death is prominent in areas such as the basal forebrain, amygdala, hippocampus, and cortical area. Over time, memory and cognitive function decline in patients thus causing dementia and eventually death [3–5]. Currently the only positive treatment is acetylcholinesterase inhibitors that merely relieve pain but not cure the disease. Exposing cell death mechanisms brought by AD may reveal novel curative strategies. The amyloid

cascade theory [6] states that extracellular plaques containing  $\beta$ -amyloid ( $\text{A}\beta$ ) peptides pathologically cause AD. These  $\text{A}\beta$  plaques mature into  $\beta$ -plated sheets and fibrillise into neuritic plates, which consequently results in microglial and astrocytic activation, oxidative injury, tau aggregation and phosphorylation, culminating in neuronal loss and synaptic dysfunction resulting in dementia [6]. Studies show that an increase in  $\text{A}\beta$  peptides triggers memory deficits [7]. These peptides are derived from the bigger amyloid precursor protein (APP).

Vascular damage is another important component of AD pathology. Degradation of the neurovascular unit (NVU) is characteristic of many neurological diseases [8]. In addition, AD vascular risk factors such as hypoglycemia, hypertension, etc. cause BBB dysfunction and damage the NVU during the process of aging [9,10]. The dysregulation of NVU leads to degeneration of nerve endings and retrograde death of cholinergic neurons [8]. This can also hinder the BBB functions, weakening the BBB's ability to clear  $\text{A}\beta$ . The  $\text{A}\beta$  then accumulates within the brain, causing chronic inflammation and further damage to the NVU [8].

### Traumatic brain injury and vascular damage

BBB leakage and vascular breakdown, including NVU impairment, have been recognized in acute brain injuries. Yearly, an estimated 235,000 Americans are hospitalized for non-fatal traumatic brain injury (TBI), 1.1 million are treated in emergency, and 50,000 die as a result of the injury [11]. Additionally, it is estimated

\* Corresponding authors. Address: 12901 Bruce B. Downs Blvd, Tampa, FL 33612, USA. Tel.: +1 813 974 3988; fax: +1 813 974 3078.

E-mail address: [cborlong@health.usf.edu](mailto:cborlong@health.usf.edu) (C.V. Borlongan).

that 43.1% of patients discharged from hospitals with acute TBI suffer from TBI-related long-term debilitation [12]. TBI is brain damage typically caused by a violent impact, blow, or jolt that results in the brain striking the inside of the skull [13]. A head puncture can also lead to TBI if the object reaches brain tissue [13]. Improvised explosive devices (IED) are often responsible for war-related TBI [14,15] because when a frontal blast wave encounters the head, the shockwave is transferred through skull, cerebrospinal fluid (CSF), and tissue. The shockwave creates negative pressure at the countercoup and has the potential to cause cavitation [16]. In addition to the current TBI target population being war fighters and veterans, TBI is also common in our daily lives, such as vehicular accidents, elderly falls, and baby accidents.

TBI has a wide range of symptoms that vary with the severity of injury. Mild TBI is typically characterized by headache, confusion, lightheadedness, dizziness, blurred vision, ringing in the ears, bad taste in the mouth, fatigue, altered sleep patterns, and behavioral or mood changes [13]. In more severe TBI cases, symptoms may be more exaggerated. For example, more severe TBI is marked by repeated vomiting or nausea, inability to awaken from sleep, seizures, pupil dilation, slurred speech, coordination loss, and confusion [13]. Many patients suffer from TBI's debilitating symptoms for the rest of their lives.


Although widely considered as an acute brain disease, TBI has been shown to be associated with chronic secondary cell death mediated by many cell death mechanisms including excitotoxicity, free radical formation, disrupted metabolism, and brain swelling [17]. Following the initial trauma, TBI continues to impair the brain by secondary neurodegeneration [17]. Therefore, it is imperative that strategies are developed to combat secondary brain tissue damage and thus limit the impact of TBI.

A major exacerbating neurodegenerative factor accompanying TBI involves a robust inflammatory reaction after the initial injury occurs [18]. What remains highly controversial is that blood-borne white cells serve as a casual factor in secondary brain injury [18]. Polymorphonuclear leukocytes and lymphocytes penetrate brain parenchyma within an hour of the primary brain insult [19]. Once the leukocytes have infiltrated the brain parenchyma, they can increase the vascular permeability in addition to releasing free radicals, activating proteases, altering the cerebral blood flow, and/or producing deleterious pro-inflammatory chemokines and cytokines [20–22].

#### **Hypothesis: vascular damage is a common pathological denominator in Alzheimer's disease and traumatic brain injury**

With the breach in vascular integrity, BBB breakdown follows TBI. Compelling evidence also shows that BBB leakage can arise secondarily to the abnormal brain activity (i.e., as seen with AD pathology) [21]. Therefore, a compromised BBB after TBI can last for years after an acute event [23]. This BBB breakdown is directly associated with an increase in the number of endothelial caveolae [24], resulting in transcytosis of plasma proteins [25,26] and a decrease in tight junction proteins [27,28]. Additionally, the BBB leakage is regarded as the arbitrator of neuroinflammatory responses, which are known to be a key pathological consequence of myriad brain pathologies including stroke, TBI, and AD [29–31].

Although the role of the BBB in the disease pathogenesis is still a developing interest of study, TBI can strongly increase the risk of developing AD [32,33]. TBI has been found to increase A $\beta$  secretion in the brain [34] and, as discussed previously, the accumulation of A $\beta$  in the brain can trigger AD [35].



**Fig. 1.** TBI-accelerated A $\beta$  plaque build-up is accompanied by vascular damage. Under normal conditions, healthy blood vessels and amyloid precursor proteins can be seen in the brain (A). Either aging of the brain or TBI or even both can cause enzymatic alteration of the APP molecule (B), resulting in aberrant aggregation of A $\beta$  plaques (C). Accumulation of A $\beta$  plaques is accompanied by vascular damage (D), which may act as a precursor to AD, as well as an exacerbating factor to TBI.

Download English Version:

<https://daneshyari.com/en/article/5812148>

Download Persian Version:

<https://daneshyari.com/article/5812148>

[Daneshyari.com](https://daneshyari.com)