



Effects of some organic pollutants on the exopolysaccharides (EPSs) produced by some *Pseudomonas* spp. strains

Dilsad Onbasli^{a,*}, Belma Aslim^b

^a Kastamonu University, Faculty of Science and Arts, Department of Biology, 37100 Kastamonu, Turkey

^b Gazi University, Faculty of Science and Arts, Department of Biology, 06500 Teknikokullar, Ankara, Turkey

ARTICLE INFO

Article history:

Received 19 November 2008

Received in revised form 29 January 2009

Accepted 29 January 2009

Available online 7 February 2009

Keywords:

Pseudomonas spp.

Exopolysaccharides (EPSs)

Some organic compounds

ABSTRACT

In this study, isolation and characterization of exopolysaccharides produced by *Pseudomonas aeruginosa* B1, *P. fluorescens* B5, *P. stutzeri* B11 and *P. putida* B15 which had been seen to produce exopolymers of potential interest in biotechnological applications were examined. To initiate the observation of the organic pollutants–polymer interactions, the yield and properties of their extracellular polysaccharide were researched. The exopolysaccharide production by these strains during growth in nutrient broth medium (control) was 41–75 mg L⁻¹. Also, *P. aeruginosa* B1, *P. fluorescens* B5, *P. stutzeri* B11 and *P. putida* B15 had exhibited high production of EPSs in presence of various organic pollutants (2,4-D, benzene, BTX and gasoline, respectively) in mineral salt medium (MSM) as a sole carbon source. EPS production by the 4 strains ranged from 40 mg L⁻¹ to 8 mg L⁻¹. Monosaccharide composition of EPS produced by these cultures were analyzed by HPLC. Results indicated that EPSs of strains contained neutral sugars and acetylated amino sugars. The neutral sugars in the EPS were mainly composed of glucose, arabinose, glycerol, ribose. The presence of galactonic acid, *N*-acetyl-D-galactosamin and *N*-acetyl-D-glucosamine indicated the acidic nature of the polysaccharide. Glycerol was the basic structural unit of EPS produced by the strains except *P. stutzeri* B11 (MSM with 1% BTX). Strain B1 (in NB medium) was found to be composed of neutral sugars (100%) while strain B1 [in MSM medium with 0.2% (v/v) 2,4-D] contained neutral sugars (70.0%), acetylated amino sugars (30.0%). Also, EPS content of strain B5 (in the NB medium) was neutral sugars (99.8%), acetylated amino sugars (0.2%) while the strain B5 [in MSM medium containing the 1% (v/v) benzene] was found to contain neutral sugars (99.9%), acetylated amino sugars (0.1%). However, EPS monomer composition by strain B11 was detected as neutral sugars (99.77%), acetylated amino sugars (0.23%) in NB medium while the strain B11 [in MSM medium with 1% (v/v) BTX] contained neutral sugars (98.2%) and acetylated amino sugars (1.8%). Lastly, in NB medium by strain B15 was found to contain neutral sugars (99.9%) and acetylated amino sugars (0.1%) while in MSM medium in the presence of 1% (v/v) gasoline it was found to contain neutral sugars (83.6%), acetylated amino sugars (16.4%). Monomer composition of control EPSs changed to different structures in the presence of various organic pollutants. Diversities of organic compounds as carbon source affected the monomer composition of EPS produced by some *Pseudomonas* spp. cultures.

© 2009 Elsevier B.V. All rights reserved.

1. Introduction

Exopolysaccharides (EPSs) are produced by many bacteria from clinical and environmental habitats and have an important function in the degradation of toxic pollutants. Recently, the pollution of soil and water by industrial chemicals is a serious problem afflicting the modern world. The use of bioremediation technologies for removing these contaminants provides a safe and economic alternative to commonly used physical–chemical treatment [1–4]. Bacteria are the primary agents of the removal/degradation of envi-

ronmental toxicants such as herbicides, pesticides, insecticides, various petrol hydrocarbons [5] and they are able to use some of these compounds as the only sources of carbon and energy. Today many microorganisms, specially *Pseudomonas* sp. are used to degrade organic pollutants in order to minimize contamination caused by several industrial activities [6]. However, in recent studies extracellular polysaccharides produced by *Pseudomonas* species have been widely reported [7–10] and polysaccharide production is a common property among these bacteria. Polysaccharides are believed to protect bacterial cells from desiccation, heavy metals, organic compounds or other environmental stresses, including host-immune responses, and to produce biofilms, thus to enhance the chances of the cells to colonize special ecological niches [11–14]. Besides this, the role of polysaccharides in industry is based on their

* Corresponding author. Tel.: +90 366 2154924; fax: +90 366 2154969.
E-mail address: donbasili@kastamonu.edu.tr (D. Onbasli).

performance as speciality chemicals, namely on their capacity to alter the basic properties of water and their propensity for emulsification, suspension, stabilization, flocculation, properties which make these polymers suitable for a specific market or application. [15]. Other important applications of microbially produced exopolymers are in the food, pharmaceutical and petroleum industries and in medical fields and they are widely accepted products of biotechnology [16].

In the present paper, it is aimed to investigate the effects of media and various organic pollutants on the yield and properties of EPS produced by *Pseudomonas aeruginosa* B1, *P. fluorescens* B5, *P. stutzeri* B11 and *P. putida* B15 strains.

2. Materials and methods

2.1. Test chemicals

The herbicide, 2,4-dichlorophenoxyacetic acid (CAS RN: 94-75-7, 98% purity) and HPLC-grade solvents were purchased from Sigma. All chemicals used for media preparation and petrol hydrocarbons [Benzene, BTX (Benzene, Toluene, Xylene) and gasoline] were ensured from Merck.

2.2. Bacterial strains

P. aeruginosa B1, *P. fluorescens* B5, *P. stutzeri* B11 and *P. putida* B15 strains used in this research were selected from between 20 cultures exhibited the highest substantial growth in presence of environmental toxicants such as herbicide (2,4-D) and various petrol hydrocarbons (benzene, BTX and gasoline) as a sole carbon source. All the four strains of *Pseudomonas* spp. were obtained from the culture collection of the Biotechnology Laboratory of Gazi University, Department of Biology Faculty of Arts and Science, in TURKEY. All of the strains were stored on Nutrient Agar Medium (Oxoid) slopes at 4 °C and stock cultures were maintained at –20 °C in 0.5% (v/v) glycerol.

2.3. Media and growth conditions

Samples were maintained in 250 mL flasks containing 50 mL of nutrient broth (NB) culture medium [17] (in g/L: peptone 2.5; NaCl 2.5; yeast extract 1.0; beef extract 0.5; pH 7.0) and mineral salt medium (MSM) [17] (in g/L: Na₂HPO₄ 4.0; KH₂PO₄ 1.5; NH₄Cl 1.0; MgSO₄·7H₂O 0.2; C₆H₈O₇·FeNH₃ 0.05; modified Hoagland trace element solution g/3,6 L: BH₃ 11.0; MnCl₂·4H₂O 7.0; AlCl₃ 1.0; CoCl₂ 1.0; CuCl₂ 1.0; KI 1.0; NiCl₂ 1.0; ZnCl₂ 1.0; BaCl₂ 0.5; KBr 0.5; LiCl 0.5; Na₂MoO₄ 0.5; SeCl₄ 0.5; SnCl₂·2H₂O 0.5; NaVO₃·H₂O 0.1; pH 7.0). NB was used as a growth medium (control) to monitor the growth of EPS production by strains while the MSM containing 1% benzene, BTX, gasoline and 0.2% 2,4-D herbicide was used to determine EPS production as well as the growth (degradation ability) of *Pseudomonas* spp.

1 mL of the active cultures was adjusted to Macfarland 5 for whole assays and inoculated inside the media. The erlenmeyer flasks were incubated at 37 °C, by using an incubator shaker (MINI-TRON) at 100–150 rpm.

NB medium was autoclaved for 15 min at 120 °C but mineral salt solutions were autoclaved separately to prevent precipitation reactions. The organic chemicals were sterilized through 0.45 µm-pore-size type HA membrane filters (Millipore Corp., Bedford, MA).

The experiment was carried out until the culture reached the stationary growth phase (72 h).

2.4. Isolation and quantification of EPS

EPS was extracted by the modified procedure of Cérantola, Bounéry, Segonds, Marty, and Montrozier [18]. Cells were harvested

at room temperature by centrifugation at 10,000 × g for 10 min. Supernatant was removed. After pellet was dissolved in 1 mL deionized distilled water, it was boiled for 15 min at 100 °C. It was then kept at room temperature for 10 min and added to 3 µL of 85% trichloroacetic acid solution (TCA). The mixture was centrifuged at 10,000 × g for 30 min. The supernatant which contained EPS was pooled and equal volume of ethanol was added. The mixture was kept at 4 °C overnight and centrifuged at 10,000 × g for 30 min again. Precipitate was then washed two times using 95% ethanol and centrifuged at 10,000 × g for 30 min. Final precipitate was dissolved in a 1-mL deionized distilled water and stored at –20 °C. Total EPS (expressed as mg per liter) was estimated in each sample by phenol-sulphuric method [19] using glucose as standard [20]. The main values were calculated from the data obtained with duplicate trials.

2.5. Characterization of EPS

The monosaccharide composition of freeze-dried exopolysaccharides samples was determined with HPLC (VARIAN ProStar) by using Metacarb 87H column (300 mm × 7.8 mm, Cat. No. 5210). The organic acids were determined with PDA detector (VARIAN 330) (210 nm), while, the egzopolysaccharides were determined with RI detector (VARIAN 350), connected following to PDA detector. The analyses conditions are; mobile phase 0.008N H₂SO₄, flow rate 0.4 mL min^{–1} and 35 °C. The analyses were accomplished by the Middle East Technical University, Central Laboratory, Molecular Biology and Biotechnology R&D Center (Ankara, Turkey) and studied as two replications.

2.6. Experimental design and statistical analysis

The experiment was performed in a completely randomized fashion with two replicates. Each analysis was done on two samples from each replicate. Results of each representative experiment were analyzed by Pearson correlation (SPSS-11). Person's correlation was used for determine any significant difference between control group of the strains and EPS production amount of the strains exposed to organic pollutants. Significance was determined at α = 0.01 level. If significant differences were indicated among treatment means ($P \leq 0.01$), means were differentiated using the least square mean test at α = 0.01.

3. Results and discussion

3.1. Isolation and quantitative determination of EPS

In this study, several organic pollutants endangering human health and environment in Turkey were used for EPS production. These organic pollutants, such as 2,4-D, benzene, BTX and gasoline, are preferred as carbon sources for the production of EPS. Hence, firstly, isolation and production of the exopolysaccharides produced by the organic pollutants-degrading bacteria were determined. The composition of the medium plays an important role in the production of EPS [21]. The organic pollutants-degrading bacteria, *P. aeruginosa* B1, *P. fluorescens* B5, *P. stutzeri* B11 and *P. putida* B15 were inoculated in both the NB (as control) and MSM medium containing 2,4-D, benzene, BTX and gasoline. EPS production by these strains ranged from 75 mg L^{–1} to 41 mg L^{–1}. In NB medium, EPS production by *P. aeruginosa* B1 (75 mg L^{–1}), *P. putida* B15 (67 mg L^{–1}), and *P. fluorescens* B5 (63 mg L^{–1}) strains was higher than that of *P. stutzeri* B11 (41 mg L^{–1}) strain. On the other hand, when these cultures were grown in medium containing organic pollutants, the EPS production was 8–40 mg L^{–1} and results were lower than that of NB medium (Table 1). EPS production of the strains exposed to organic pollutants were significantly decreased according to

Download English Version:

<https://daneshyari.com/en/article/581308>

Download Persian Version:

<https://daneshyari.com/article/581308>

[Daneshyari.com](https://daneshyari.com)