Accepted Manuscript



Phosphodiesterase 10A inhibitor, MP-10 (PF-2545920), produces greater induction of c-Fos in dopamine D2 neurons than in D1 neurons in the neostriatum

Jonathan M. Wilson, Ann Marie L. Ogden, Sally Loomis, Gary Gilmour, Anthony J. Baucum, II, Teri L. Belecky-Adams, Kalpana M. Merchant

PII: S0028-3908(15)30054-X

DOI: 10.1016/j.neuropharm.2015.08.008

Reference: NP 5954

To appear in: *Neuropharmacology*

Received Date: 11 June 2015

Revised Date: 31 July 2015

Accepted Date: 4 August 2015

Please cite this article as: Wilson, J.M., Ogden, A.M.L., Loomis, S., Gilmour, G., Baucum II., A.J., Belecky-Adams, T.L., Merchant, K.M., Phosphodiesterase 10A inhibitor, MP-10 (PF-2545920), produces greater induction of c-Fos in dopamine D2 neurons than in D1 neurons in the neostriatum, *Neuropharmacology* (2015), doi: 10.1016/j.neuropharm.2015.08.008.

Phosphodiesterase 10A inhibitor, MP-10 (PF-2545920), produces greater induction of c-Fos in dopamine D2 neurons than in D1 neurons in the neostriatum

Jonathan M. Wilson[a,c,*], Ann Marie L. Ogden[a], Sally Loomis[b], Gary Gilmour[b], Anthony J. Baucum II[c,d], Teri L. Belecky-Adams[c] and Kalpana M. Merchant[c]

[a]Lilly Research Laboratories, Eli Lilly and Company, Indianapolis, Indiana, 46225, USA; [b]Erl Wood Manor, Sunninghill Road, Windlesham, Surrey GU20 6PH, England, UK; [c]Department of Biology, Center for Developmental and Regenerative Biology, [d]Stark Neurosciences Research Institute, Indiana University-Purdue University Indianapolis 723 West Michigan Street, Indianapolis, Indiana, 46202, USA

*Corresponding Author:

Jonathan M. Wilson

Eli Lilly and Company

Indianapolis, IN 46225

Telephone Number: (317) 651-3810

Fax Number: (317) 277-4499

Email Address: wilsonjo@lilly.com