

Cuprizone decreases intermediate and late-stage progenitor cells in hippocampal neurogenesis of rats in a framework of 28-day oral dose toxicity study

Hajime Abe, Takeshi Tanaka, Masayuki Kimura, Sayaka Mizukami, Fumiyo Saito, Nobuya Imatanaka, Yumi Akahori, Toshinori Yoshida, Makoto Shibutani

PII: S0041-008X(15)30012-0
DOI: doi: 10.1016/j.taap.2015.06.005
Reference: YTAAP 13395

To appear in: *Toxicology and Applied Pharmacology*

Received date: 13 March 2015
Revised date: 31 May 2015
Accepted date: 4 June 2015

Please cite this article as: Abe, Hajime, Tanaka, Takeshi, Kimura, Masayuki, Mizukami, Sayaka, Saito, Fumiyo, Imatanaka, Nobuya, Akahori, Yumi, Yoshida, Toshinori, Shibutani, Makoto, Cuprizone decreases intermediate and late-stage progenitor cells in hippocampal neurogenesis of rats in a framework of 28-day oral dose toxicity study, *Toxicology and Applied Pharmacology* (2015), doi: 10.1016/j.taap.2015.06.005

Cuprizone decreases intermediate and late-stage progenitor cells in hippocampal neurogenesis of rats in a framework of 28-day oral dose toxicity study

Hajime Abe[a,b], Takeshi Tanaka[a,b], Masayuki Kimura[a,b], Sayaka Mizukami[a,b], Fumiyo Saito[c], Nobuya Imatanaka[c], Yumi Akahori[c], Toshinori Yoshida[a], and Makoto Shibutani[a,*]

[a]Laboratory of Veterinary Pathology, Tokyo University of Agriculture and Technology, 3-5-8 Saiwai-cho, Fuchu-shi, Tokyo 183-8509, Japan

[b]Pathogenetic Veterinary Science, United Graduate School of Veterinary Sciences, Gifu University, 1-1 Yanagido, Gifu-shi, Gifu 501-1193, Japan

[c]Chemicals Evaluation and Research Institute, Japan, 1-4-25 Koraku, Bunkyo-ku, Tokyo 112-0004, Japan

Corresponding author: Prof. Makoto Shibutani, D.V.M., Ph.D.

Tel/Fax: +81-42-367-5771; E-mail: mshibuta@cc.tuat.ac.jp