



## Cardio protective effect of *Coriandrum sativum* L. on isoproterenol induced myocardial necrosis in rats

Dipak K. Patel<sup>a</sup>, Swati N. Desai<sup>a</sup>, Hardik P. Gandhi<sup>b</sup>, Ranjitsinh V. Devkar<sup>a,\*</sup>, A.V. Ramachandran<sup>a</sup>

<sup>a</sup>Division of Phytotherapeutics and Metabolic Endocrinology, Department of Zoology, Faculty of Science, The Maharaja Sayajirao University of Baroda, Vadodara-390002, Gujarat, India

<sup>b</sup>Department of Pharmacy, Faculty of Technology and Engineering, The Maharaja Sayajirao University of Baroda, Vadodara-390002, Gujarat, India

### ARTICLE INFO

#### Article history:

Received 22 August 2011

Accepted 15 June 2012

Available online 28 June 2012

#### Keywords:

*Coriandrum sativum* L.

Seed

Methanolic extract

Antioxidants

Isoproterenol

Cardiotoxicity

### ABSTRACT

The preventive effect of *Coriandrum sativum* L. (CS) on cardiac damage was evaluated by Isoproterenol (IP) induced cardiotoxicity model in male *Wistar* rats. Rats were pretreated with methanolic extract of CS seeds at a dose of 100, 200 or 300 mg/kg orally for 30 days and they were subsequently administered (s.c.) with IP (85 mg/kg body weight) for the last two days. IP treated rats showed increased LPO, decreased levels of endogenous antioxidants and ATPases in the cardiac tissue together with increased plasma lipids and markers of cardiac damage. TTC staining showed increased infarct areas while HXE staining showed myofibrillar hypertrophy and disruption. CS (200 and 300 mg/kg body weight) pretreatment significantly prevented or resisted all these changes. Our results show that methanolic extract of CS is able to prevent myocardial infarction by inhibiting myofibrillar damage. It is also concluded that, the rich polyphenolic content of CS extract is responsible for preventing oxidative damage by effectively scavenging the IP generated ROS.

© 2012 Elsevier Ltd. All rights reserved.

### 1. Introduction

Epidemiological studies predict an ominous prevalence of cardio vascular diseases globally as well as in India during next decade (Lopez and Murray, 1998; Gilski and Borkenhagen, 2005). Myocardial infarction, a highly prevalent ischemic condition characterized by tissue necrosis develops essentially due to an imbalance between oxygen need and actual supply (De Bono and Boon, 1992) and results in irreversible histopathological damages and subsequent cardiovascular complications (Gross and Auchampach, 2007).

Isoproterenol (IP), a synthetic catecholamine and  $\beta$ -adrenergic agonist increases heart rate and exhaust energy reservoir of cardiac myocytes leading to cell death. It induces myocardial necrosis via multiple modes of action in experimental animals. It is essentially manifest by its stimulation of sarcolemmal adenylate cyclase and  $\text{Na}^+$  and  $\text{Ca}^{2+}$  channels resulting in exaggerated influx of  $\text{Ca}^{2+}$  and energy consumption and consequent cell death (Milei et al., 1978). Free radicals produced by IP initiate the peroxidation of membrane bound polyunsaturated fatty acids (PUFAs) leading to both structural and functional myocardial injury (Thompson and Hess, 1986). IP-induced myocardial necrosis serves as an excellent experimental model to study catecholamines induced cardiac

dysfunction and also to evaluate the possible cardioprotective efficacy of various natural and synthetic agents.

*Coriandrum sativum* L. (Apiaceae) (CS) is an ubiquitous annual herb, the leaves and seeds of which form a key ingredient of Middle Eastern, Mediterranean, Indian, Latin American, African and South-east Asian cuisines. Apart from its usage as a condiment, decoction and tincture of powdered seeds of CS find usage either alone or in combination with other herbals in the treatment of cough, dysentery, sore throat, convulsion, insomnia and anxiety (Grieve, 1971). An extract of CS seeds is also reported to have therapeutic potential against diabetes, cardiovascular and urinary disorders (Eguale et al., 2007; Emamghoreishi et al., 2005). Phytochemical analysis of CS seeds has revealed the presence of polyphenols (rutin, ferulic acid, gallic acid, chlorogenic acid and caffeic acid derivatives), flavonoids (quercetin and isoquercetin) and  $\beta$ -carotenoids (Melo et al., 2003). The oil of CS seeds is rich in  $\alpha$  and  $\beta$ -pinene, camphor, citronellol, coriandrol, *p*-cymene, geraniol, geranyl acetate, limonene, linalool, myrcene,  $\alpha$  and  $\beta$  phellandrene and terpinene besides many water soluble compounds such as monoterpene glycosides and their derivatives (Sergeeva, 1975; Ishikawa et al., 2003). The reported pharmacological actions of CS are many with its oil shown to possess antifungal (Garg and Siddiqui, 1992) and antimicrobial (Baratta et al., 1998) properties and seed extract shown to possess hypoglycemic (Gray and Flatt, 1999), hypolipidemic (Chithra and Leelamma, 1997; Chithra and Leelamma, 1999; Lal et al., 2004), hypocholesterolemic (Dhanapakiam et al., 2008),

\* Corresponding author. Tel.: +91 9825935445; fax: +91 0265 2342109.

E-mail address: [phyto\\_met@yahoo.com](mailto:phyto_met@yahoo.com) (R.V. Devkar).

anti-insulin resistance activity (Patel et al., 2011), antihypertensive (Medhin et al., 1986) and antioxidant (Melo et al., 2003; Ramadan et al., 2003; Bajpai et al., 2005) competence.

Several pre-clinical and clinical studies involving pretreatment with vitamins and antioxidants have demonstrated their potential to prevent myocardial damage (Singh et al., 1994; Senthil et al., 2004). Previously Hashim et al. (2005) have investigated that hydro-methanolic extract of CS seed had strong antioxidant property and it had prevented oxidative damage induced by H<sub>2</sub>O<sub>2</sub> to lymphocytes. The present study was designed to assess cardioprotective potential of hydro-methanolic extract of the customarily used spice CS seeds in IP induced multifocal myocardial necrosis in rats.

## 2. Materials and methods

### 2.1. Plant material and preparation of extract

CS plants were collected in the seedling months (February and March) and Dr. P.S. Nagar, Department of Botany, The M.S. University of Baroda identified the plant and a sample specimen was deposited in the herbarium of the Department of Botany. Hundred grams of powdered dry seeds soaked in methanol:water (80:20 v/v) at room temperature was allowed to stand for seven days. Resultant extract filtered through a muslin cloth was concentrated in a rotary evaporator under reduced pressure to obtain a thick semisolid brown paste (Qaiser et al., 2009). The final yield obtained was 8.3 g (w/w).

### 2.2. Experimental animals

Adult male Wistar rats (150–200 gm; obtained from Zydus Cadila Research Centre, Ahmedabad, Gujarat, India) were housed under standard animal house conditions (23 ± 2 °C; LD 12:12 and 45–50% humidity) and provided with pelleted diet (M/S Pranav agro, Ltd., Baroda, India) and water *ad libitum*. The animals were maintained as per the guidelines of the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA) India and the experimental protocol approved by the animal ethical committee of the Department of Zoology, The M.S. University of Baroda, Vadodara (Approval No.827/ac/04/CPCSEA).

### 2.3. Experiment design

Thirty animals were randomly divided into five groups of six animals each. Group I (NC) served as control and received 0.5% Carboxy methyl cellulose (CMC; *p.o.*) for 28 days and normal saline (*s.c.*) on days 29 and 30. Group II (IP) served as positive control rats and received 0.5 CMC (*p.o.*) for 28 days and isoproterenol (85 mg/kg body weight, *s.c.*) on days 29 and 30 while, the remaining groups [Group III (IP + CS100), Group IV (IP + CS200) and group V (IP + CS300)] received respectively 100, 200 and 300 mg/kg body weight of CS extract daily for 28 days (*p.o.*) and IP (85 mg/kg, *s.c.*) on days 29 and 30. The protocol for IP treatment schedule was as per the previous works from this laboratory (Jadeja et al., 2010; Thounaojam et al., 2011). At the end of the experimental period (i.e. 31st day), animals were fasted overnight (12 h) and blood samples were collected from retro-orbital sinus under mild ether anesthesia. Plasma was obtained by cold centrifugation of samples at 3000 rpm for 10 min. Later, animals were sacrificed by cervical dislocation under mild anesthesia and heart was excised and stored at –80 °C for further evaluations. A piece of cardiac tissue was fixed in 10% paraformaldehyde for paraffin wax histology.

### 2.4. Plasma markers of cardiac damage

Plasma levels of creatine phosphokinase- MB (CK-MB), lactate dehydrogenase (LDH), aspartate transaminase (AST), alanine transaminase (ALT) and uric acid were ascertained by using commercially available kits (Reckon Diagnostic Ltd., Vadodara, India).

**Table 1**

Effect of CS seed extract on plasma markers of cardiac damage.

Parameters	NC	IP	IP + CS100	IP + CS200	IP + CS300
CkMB <sup>§</sup>	75.66 ± 6.91	218.20 ± 29.16 <sup>c</sup>	171.20 ± 9.19 <sup>B</sup>	133.10 ± 6.16 <sup>C</sup>	80.10 ± 9.03 <sup>C</sup>
LDH <sup>#</sup>	82.71 ± 6.50	189.60 ± 7.36 <sup>c</sup>	149.70 ± 4.32 <sup>C</sup>	126.00 ± 4.15 <sup>C</sup>	85.60 ± 6.35 <sup>C</sup>
AST <sup>*</sup>	30.33 ± 1.99	61.17 ± 2.24 <sup>c</sup>	50.33 ± 1.76 <sup>B</sup>	43.50 ± 1.91 <sup>C</sup>	31.67 ± 1.02 <sup>C</sup>
ALT <sup>*</sup>	19.33 ± 1.11	44.83 ± 2.18 <sup>c</sup>	36.67 ± 1.82 <sup>A</sup>	31.00 ± 1.73 <sup>C</sup>	22.17 ± 1.99 <sup>C</sup>
Uric acid <sup>@</sup>	1.91 ± 0.21	7.01 ± 0.47 <sup>c</sup>	5.24 ± 0.41 <sup>A</sup>	3.72 ± 0.19 <sup>C</sup>	2.14 ± 0.21 <sup>C</sup>

Where, § = IU/l, # = U/l, \* = KA Units/l, @ = mg/dl. n = 6. Data were expressed as mean ± S.E.M. a (*p* < 0.05), b (*p* < 0.01), c (*p* < 0.001) when NC vs. IP and A (*p* < 0.05), B (*p* < 0.01), C (*p* < 0.001) when IP vs. IP + CS.

### 2.5. Plasma lipid profile

Triglyceride (TG), total cholesterol (TC) and high density lipoprotein (HDL) content were assayed by using commercially available kits (Recon Diagnostic, Ltd., Vadodara, India). Lowdensity lipoprotein (LDL) and Very low-density lipoprotein (VLDL) were calculated by Friedewald's formula (Friedewald et al., 1972).

### 2.6. Cardiac antioxidants and Lipid peroxidation (LPO)

Cardiac tissue from control and treated groups was weighed and homogenized (10%w/v) in chilled Tris buffer (10 mM; pH 7.4) and centrifuged at 10,000 g for 20 min at 0 °C. Clear supernatant was used to assay superoxide dismutase (SOD; Marklund and Marklund, 1974), catalase (CAT; Aebi, 1983), glutathione peroxidase (GPx; Rotruck et al., 1973), glutathione s-transferase (GST; Habig et al., 1974), reduced glutathione (GSH; Beutler, 1963), vitamin E (Vit. E; Baker and Frank, 1968), total protein content (Lowry et al., 1951) and lipid peroxidation levels (LPO; Buege and Aust, 1978). Total ascorbic acid content (AA) was measured as per Roe and Kuehler (1943) by preparing homogenates of fresh cardiac tissue in 6% Trichloro acetic acid.

### 2.7. Cardiac ATPases

Pellets obtained from tissue homogenate after centrifugation was re-suspended in ice-cold Tris buffer (10 mM, pH 7.4) to get a final concentration of 10% and was used for the estimation of Na<sup>+</sup> K<sup>+</sup> ATPase (Bonting et al., 1970), Ca<sup>2+</sup> ATPase (Hjerken and Pan, 1983) and Mg<sup>2+</sup>ATPase (Ohinishi et al., 1982). Protein was estimated according to the method of Lowry et al. (1951).

### 2.8. Macroscopic and microscopic evaluation of cardiac tissue

Heart tissue slices (approx. 2–3 mm thick) transversely cut across the ventricle were kept in a covered glass dish containing 1% TTC (2, 3, 5- triphenyltetrazolium chloride; Sigma, St. Louis, MO) solution and incubated at 37 °C for 20 min for differentiation of viable tissue from necrotic areas (Li et al., 2011).

Heart samples from control and treated rats were fixed in 4% buffered paraformaldehyde, dehydrated in graded alcohol series and embedded in paraffin wax. Five micrometer thick sections cut (by Leica RM2155 Microtome) and stained with haematoxylin-eosin, were photographed with Canon power shot S72 digital Camera (200×) attached to a Leica microscope.

### 2.9. Statistical analysis

Statistical analysis of data was done by one way ANOVA followed by Bonferroni's multiple comparison test and results were expressed as mean ± S.E.M (Using Graph Pad Prism version 3.0 for Windows, Graph Pad Software, San Diego California USA).

## 3. Results

### 3.1. Plasma markers of cardiac damage

IP treated rats showed significant (*p* < 0.005) increment in the plasma levels of CK-MB, LDH, AST, ALT and uric acid compared to NC rats. Pretreatment of IP rats with CS prevented the IP induced increase in the serum levels of these parameters in a dose dependent manner (Table 1).

### 3.2. Plasma lipid profile

IP treatment recorded significant (*p* < 0.005) increase in plasma TG, TC, LDL, and VLDL and decrement in HDL levels compared to the NC group. CS treatment showed dose dependent decrement

Download English Version:

<https://daneshyari.com/en/article/5852208>

Download Persian Version:

<https://daneshyari.com/article/5852208>

[Daneshyari.com](https://daneshyari.com)