Accepted Manuscript

Mechanisms of Exercise-Induced Hypoalgesia

Kelli F. Koltyn, Angelique G. Brellenthin, Dane B. Cook, Nalini Sehgal, Cecilia Hillard

PII: S1526-5900(14)00915-8

DOI: 10.1016/j.jpain.2014.09.006

Reference: YJPAI 2986

To appear in: *Journal of Pain*

Received Date: 28 February 2014

Revised Date: 17 July 2014

Accepted Date: 12 September 2014

Please cite this article as: Koltyn KF, Brellenthin AG, Cook DB, Sehgal N, Hillard C, Mechanisms of Exercise-Induced Hypoalgesia, *Journal of Pain* (2014), doi: 10.1016/j.jpain.2014.09.006.

Mechanisms of Exercise-Induced Hypoalgesia

Kelli F. Koltyn[1], Angelique G. Brellenthin,[1] Dane B. Cook,[1] Nalini Sehgal,[2] and Cecilia Hillard[3]

Department of Kinesiology[1] and Department of Rehabilitation Medicine[2]
University of Wisconsin-Madison[1,2]
Department of Pharmacology and Toxicology[3]
Medical College of Wisconsin[3]

Research was conducted at the University of Wisconsin Hospital and Clinics: Clinical Research Unit

Correspondence should be sent to:

Kelli F. Koltyn, Ph.D.
2000 Observatory Drive
University of Wisconsin-Madison
Madison, WI 53706-1121
Koltyn@education.wisc.edu
Phone: (608) 262-4234
Fax: (608) 262-1656

Text pages including Figures and Tables: 33
Figures: 5 and Tables: 1