

Lung Cancer Hormesis in High Impact States Where Nuclear Testing Occurred

Steven Lehrer, Kenneth E. Rosenzweig

Abstract

Hormesis is a favorable biological response to low toxin exposure. In the case of radiation, large doses are carcinogenic, but low doses might be protective. Lung cancer incidence is significantly lower in states affected by nuclear testing. Our analysis adds to the body of evidence suggesting that the linear no threshold model of radiation carcinogenicity in lung cancer might not be correct. Low-level radiation exposure might protect against lung cancer rather than cause it.

Background: Hormesis is a favorable biological response to low toxin exposure. In the case of radiation, large doses are carcinogenic, but low doses might be protective. In the current study, we analyzed lung cancer incidence in high-impact radiation states where nuclear testing occurred and compared it with lung cancer incidence in the remaining normal-impact radiation states and the District of Columbia. **Materials and Methods:** Lung cancer incidence data were from the American Cancer Society. Tobacco use 2012 data were from the Centers for Disease Control and Prevention. The distribution of states grouped according to lung cancer incidence interval was from the Centers for Disease Control and Prevention. Total background radiation measurements (terrestrial + cosmic + radon) were from Assessment of Variations in Radiation Exposure in the United States (2005). Data on high- and normal-impact states were from the National Radiation Exposure Screening & Education Program (RESEP). Congress passed the Radiation Exposure Compensation Act Amendments of 2000, creating RESEP, to help thousands of people diagnosed with cancer and other diseases caused by exposure to nuclear fallout or radioactive materials such as uranium. These people live in 12 high-impact states where nuclear testing had occurred. High-impact states were not designated according to measurements of background radiation. **Results:** Lung cancer incidence is significantly lower in high-impact states in men ($t = 5.4$ for unequal variance; $P < .001$) and women ($t = 3.0$; $P < .001$). The clustering of the 12 high-impact states in the 2 lowest lung cancer incidence intervals (26.8-56.9 and 57.0-63.2) is statistically significant ($P < .001$, Fisher exact test, 2-tailed). Because cigarette smoking is ordinarily the most powerful risk factor for lung cancer, multivariate linear regression analysis of the effect of U.S. state group (normal-impact, high-impact, or extra high-impact for Nevada, Utah, and Arizona) on lung cancer incidence in men and women was performed. (In Nevada, Utah, and Arizona, men and women would have been downwind.) The U.S. state group impact was significant ($P < .001$ for men; $P = .015$ for women). The effect of percentage of smokers in the population was significant ($P < .001$ for men; $P < .001$ for women). The effect of total background radiation was significant ($P = .029$ for men; $P < .029$ for women); like the state group impact, more background radiation exposure was associated with less lung cancer. **Conclusion:** Hormesis is still mired in controversy. Yet, it is of vital medical importance because of the continuing debate over whether the low-level radiation doses from diagnostic x-ray procedures, such as computed tomography scans, are harmful. Our analysis adds to the body of evidence suggesting that the linear no threshold model of radiation carcinogenicity in lung cancer might not be correct. Low-level radiation exposure might protect against lung cancer rather than cause it.

Clinical Lung Cancer, Vol. 16, No. 2, 152-5 © 2015 Elsevier Inc. All rights reserved.

Keywords: Background, Linear no threshold model, Radiation

Department of Radiation Oncology, Icahn School of Medicine at Mount Sinai, NY

Submitted: Aug 21, 2014; Accepted: Sep 30, 2014; Epub: Oct 13, 2014

Address for correspondence: Steven Lehrer, MD, Mount Sinai Medical Center Box 1236, 1 Gustave L. Levy Place, New York, NY 10029
 Fax: 212-245-9708; e-mail contact: steven.lehrer@mssm.edu

Introduction

Hormesis is a favorable biological response to low toxin exposure. A pollutant or toxin demonstrating hormesis has the opposite effect in small doses as in large doses.¹ In the case of radiation, large doses are carcinogenic. However, Frigerio et al found lower overall cancer rates in U.S. states with high-impact radiation.²

Large doses of radiation from radon in houses, exceeding 10 pCi per liter (2000 mrem/y), are associated with increased lung cancer incidence.^{3,4} However, low-dose radon home exposure is associated with reduced rates of lung cancer. Bogen compared Environmental Protection Agency radon data, county by county, with lung cancer mortality records for women. He confirmed the inverse correlation between lung cancer and radon.⁵

Cohen examined the linear no threshold (LNT) model of radiation carcinogenicity in lung cancer.⁶ This model is used in radiation protection to quantify radiation exposure and set regulatory limits. LNT assumes that the long-term, biological damage caused by ionizing radiation (in other words, the cancer risk) is directly proportional to the dose. LNT presumes that radiation is always harmful with no safety threshold, and the sum of multiple small exposures has the same effect as 1 large exposure (ie, response linearity).⁷

Cohen found that the LNT model overstated the effects of radiation. For example, lung cancer incidence in the high-radon area of Cumberland County, Pennsylvania was lower than the Pennsylvania average.⁶ Thompson found that the maximum hormesis for lung cancer occurred at 70 Bq m³ or 350 mrem/y.⁸ Nevertheless, hormesis in lung cancer is still controversial.

People who live in 12 U.S. states where nuclear weapons testing occurred are classified as living in high-impact states. These states are Arizona, Colorado, Idaho, Nevada, New Mexico, North Dakota, Oregon, South Dakota, Texas, Utah, Washington, and Wyoming.⁹ In the current study, we analyzed lung cancer incidence in high-impact states and compared it with lung cancer incidence in the remaining normal-impact states and the District of Columbia.

Materials and Methods

Lung cancer incidence data were from the American Cancer Society.¹⁰ Tobacco use 2012 data were from the Centers for Disease Control and Prevention.¹¹ The distribution of states grouped according to lung cancer incidence interval (used in Table 1) data were from the Centers for Disease Control and Prevention.¹² Total

background radiation measurements (terrestrial + cosmic + radon) were from the Assessment of Variations in Radiation Exposure in the United States.¹³

Data on high- and normal-impact states were from the National Radiation Exposure Screening & Education Program (RESEP).⁹ Congress passed the Radiation Exposure Compensation Act Amendments of 2000, creating RESEP, to help thousands of people diagnosed with cancer and other diseases caused by exposure to nuclear fallout or nuclear materials such as uranium. RESEP set the following criteria to identify affected individuals in high-impact states:

- Uranium Mine Worker: a person who operated or otherwise worked for at least 1 year, or could establish radon exposure equivalent to 40 working level months, in above-ground or underground uranium mines in specified states (AZ, CO, ID, OR, ND, NM, SD, TX, UT, WA, WY) during the period beginning January 1, 1942 and ending December 31, 1971.
- Uranium Mill Worker: a person who was employed for at least 1 year as a uranium mill worker in specified states (AZ, CO, ID, OR, ND, NM, SD, TX, UT, WA, WY) during the period beginning January 1, 1942 and ending December 31, 1971.
- Uranium Ore Transporter: a person who was employed for at least 1 year as a transporter of uranium ore or vanadium-uranium ore from a uranium mine or uranium mill located in a specified state (AZ, CO, ID, OR, ND, NM, SD, TX, UT, WA, WY) during the period beginning January 1, 1942 and ending December 31, 1971.
- Downwinder: a person who was exposed to fallout from the atmospheric detonation of nuclear devices at the Nevada Test Site because of their physical presence in Arizona counties: Apache, Coconino, Gila, a portion of Mohave County (north of the Grand Canyon), Navajo, or Yavapai; Nevada counties: Eureka, Lander, Lincoln, Nye, White Pine, and a portion of Clark; Utah counties: Beaver, Garfield, Iron, Kane, Millard, Piute, San Juan, Sevier, Washington, or Wayne. Downwind counties were determined based on wind patterns around the dates of atmospheric nuclear tests at the Nevada Test Site. Under the current law, only portions of Nevada, Utah, and Arizona are considered downwind. The other 9 high-impact states have significant concentrations of uranium miners, millers, or ore transporters. Nevada also has a significant number of “onsite participants.” Some lawmakers have introduced bills to expand the high-impact area, most recently the Radiation Exposure Compensation Act Amendments of 2013, but the amendments have not been passed by Congress.

High-impact states and normal-impact states were not designated according to measurements of background radiation.

Results

Lung cancer incidence was significantly lower in high-impact states in men ($t = 5.4$ for unequal variance; $P < .001$) and women ($t = 3.0$; $P < .001$; Figure 1).

High- and normal-impact states grouped according to lung cancer incidence interval (cases per 100,000) are shown in Table 1. The clustering of the 12 high-impact states in the 2 lowest incidence

Table 1 High- and Normal-Impact Radiation States Grouped According to Lung Cancer Incidence Interval (Cases Per 100,000)

Incidence Interval	States
26.8 to 56.9	ARIZONA , California, COLORADO , Hawaii, IDAHO , Montana, New Jersey, NEW MEXICO , NORTH DAKOTA , OREGON , TEXAS , UTAH , and WYOMING
57.0 to 63.2	Connecticut, District of Columbia, Florida, Kansas, Maryland, Nebraska, NEVADA , New York, SOUTH DAKOTA , Virginia, WASHINGTON , and Wisconsin
63.3 to 68.4	Alaska, Delaware, Georgia, Illinois, Iowa, Massachusetts, New Hampshire, Ohio, Oklahoma, Pennsylvania, Rhode Island, and South Carolina
68.5 to 97.3	Alabama, Indiana, Kentucky, Louisiana, Maine, Michigan, Mississippi, Missouri, North Carolina, Tennessee, Vermont, and West Virginia

The 12 high-impact states are in all capital letters and bold. The clustering of the 12 high-impact states in the 2 lowest incidence intervals (26.8-56.9 and 57.0-63.2) is statistically significant ($P < .001$; Fisher exact test, 2-tailed). No data were available for Arkansas and Minnesota.

Download English Version:

<https://daneshyari.com/en/article/5882686>

Download Persian Version:

<https://daneshyari.com/article/5882686>

[Daneshyari.com](https://daneshyari.com)