


## Accepted Manuscript

Title: Prevalence of resistant hypertension and associated factors in Japanese subjects with type 2 diabetes

Author: H. Yokoyama S. Araki S. Watanabe J. Honjo S. Okizaki D. Yamada R. Shudo H. Shimizu H. Sone M. Haneda



PII: S0168-8227(15)00364-2  
DOI: <http://dx.doi.org/doi:10.1016/j.diabres.2015.08.007>  
Reference: DIAB 6457

To appear in: *Diabetes Research and Clinical Practice*

Received date: 2-3-2015  
Revised date: 4-8-2015  
Accepted date: 18-8-2015

Please cite this article as: H. Yokoyama, S. Araki, S. Watanabe, J. Honjo, S. Okizaki, D. Yamada, R. Shudo, H. Shimizu, H. Sone, M. Haneda, Prevalence of resistant hypertension and associated factors in Japanese subjects with type 2 diabetes, *Diabetes Research and Clinical Practice* (2015), <http://dx.doi.org/10.1016/j.diabres.2015.08.007>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Highlights

The prevalence of treatment resistant hypertension is unclear in diabetes. This was estimated by mean values of office blood pressure over one year. It was evaluated with a high achievement of BP goals and adequate adherence. It was associated with cardiovascular risks and familial predisposition to hypertension. Treatment resistant hypertension is common and important in diabetes.

Download English Version:

<https://daneshyari.com/en/article/5899209>

Download Persian Version:

<https://daneshyari.com/article/5899209>

[Daneshyari.com](https://daneshyari.com)